

08-14-00

A

10860 U.S. PTO

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

10862 U.S. PTO 09/637614 08/11/00

Patent Application of: Shohei Koide
Title: ARTIFICIAL ANTIBODY POLYPEPTIDES
Attorney Docket No.: 109.034US3

PATENT APPLICATION TRANSMITTAL

BOX PATENT APPLICATION

Commissioner for Patents
Washington, D.C. 20231

We are transmitting herewith the following attached items and information (as indicated with an "X"):

- X Return postcard.
X DIVISIONAL of prior Patent Application No. 09/096,749 (under 37 CFR § 1.53(b)) comprising:
X Specification (80 pgs, including claims numbered 1 through 39 and a 1 page Abstract).
X Formal Drawing(s) (15 sheets).
X Copy of signed Combined Declaration and Power of Attorney (3 pgs) from prior application.
X Incorporation by Reference: The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied herewith, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
X Check in the amount of \$345.00 to pay the filing fee.
X Information Disclosure Statement (1 pgs), Form 1449 (3 pgs). References NOT enclosed, cited in prior application.
X Preliminary Amendment (6 pgs), and redlined copy of Fig. 2 (1 pg)..
X Small Entity Statement filed in prior application; status still proper and desired.
X Communication Under 37 C.F.R. 1.821 (1 pgs), with SEQUENCE LISTING (40 pgs).

The filing fee has been calculated below as follows:

Table with 5 columns: Item, No. Filed, No. Extra, Rate, Fee. Rows include TOTAL CLAIMS, INDEPENDENT CLAIMS, MULTIPLE DEPENDENT CLAIMS PRESENTED, BASIC FEE, and TOTAL.

Please charge any additional required fees or credit overpayment to Deposit Account No. 19-0743.

SCHWEGMAN, LUNDBERG, WOESSNER & KLUTH, P.A.
P.O. Box 2938, Minneapolis, MN 55402 (612-373-6900)

By: [Signature]
Atty: Ann S. Viksnins
Reg. No. 37,748

Customer Number 21186

"Express Mail" mailing label number: EL709305975US Date of Deposit: August 11, 2000
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner for Patents, Box Patent Application, Washington, D.C. 20231.

By: Shawn L. Hise Signature: [Signature]

S/N Unknown

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:	Shohei Koide	Examiner:	Unknown
Serial No.:	Unknown	Group Art Unit:	Unknown
Filed:	Herewith	Docket:	109.034US3
Title:	ARTIFICIAL ANTIBODY POLYPEPTIDES		

PRELIMINARY AMENDMENT

BOX PATENT APPLICATION

Commissioner for Patents

Washington, D.C. 20231

Sir:

Prior to examination, please amend the above-identified divisional application as follows.

In the Specification

Please insert the enclosed SEQUENCE LISTING into the specification.

On page 1, after the title, please insert the following: --This application is a divisional of U.S. Serial No. 09/096,749, filed June 12, 1998, which claims priority under 35 U.S.C. § 119(e) to provisional application U.S. Serial No. 60/049,410, filed June 12, 1997. These applications are incorporated herein by reference.--

On page 12, line 13 after "Amino acid sequence" insert -- (SEQ ID NO:110)--.

On page 12, line 18 before ", preceding" insert -- (SEQ ID NO:114)--.

On page 13, line 5 after "sequences" insert -- (SEQ ID NO:111 and SEQ ID NO:112, respectively)--.

On page 14, line 5 after "pLB25.1" insert -- (containing SEQ ID NO:115)-- and after "pLB25.4" insert -- (containing SEQ ID NO:116)--.

On page 14, line 6 after "pLB24.1" insert -- (containing SEQ ID NO:117)-- and after "pLB24.3" insert -- (containing SEQ ID NO:118)--.

On page 18, line 16 after "(RGD)" insert -- (SEQ ID NO:113) --.

On page 21, please make the following changes:

at the end of line 9 insert -- (SEQ ID NO:1)--;

at the end of line 10 insert -- (SEQ ID NO:2)--;

at the end of line 12 insert -- (SEQ ID NO:3)--;

at the end of line 13 insert -- (SEQ ID NO:4)--;

PRELIMINARY AMENDMENT

Serial Number: Unknown
Filing Date: Herewith
Title: ARTIFICIAL ANTIBODY POLYPEPTIDES

Page 2
Dkt: 109.034US3

at the end of line 14 insert -- (SEQ ID NO:5)--;
at the end of line 15 insert -- (SEQ ID NO:6)--;
at the end of line 16 insert -- (SEQ ID NO:7)--;
at the end of line 17 insert -- (SEQ ID NO:8)--;
at the end of line 18 insert -- (SEQ ID NO:9)--;
at the end of line 19 insert -- (SEQ ID NO:10)--;
at the end of line 20 insert -- (SEQ ID NO:11)--;
at the end of line 21 insert -- (SEQ ID NO:12)--.

On page 25 please make the following changes:

at the end of line 7 insert -- (SEQ ID NO:13)--;
at the end of line 8 insert -- (SEQ ID NO:14)--;
at the end of line 9 insert -- (SEQ ID NO:15)--;
at the end of line 10 insert -- (SEQ ID NO:16)--;
at the end of line 11 insert -- (SEQ ID NO:17)--;
at the end of line 12 insert -- (SEQ ID NO:18)--;
at the end of line 13 insert -- (SEQ ID NO:19)--;
at the end of line 14 insert -- (SEQ ID NO:20)--;
at the end of line 15 insert -- (SEQ ID NO:21)--;
at the end of line 16 insert -- (SEQ ID NO:22)--;
at the end of line 17 insert -- (SEQ ID NO:23)--;
at the end of line 18 insert -- (SEQ ID NO:24)--;
at the end of line 19 insert -- (SEQ ID NO:25)--;
at the end of line 20 insert -- (SEQ ID NO:26)--;
at the end of line 21 insert -- (SEQ ID NO:27)--;
at the end of line 22 insert -- (SEQ ID NO:28)--;
at the end of line 23 insert -- (SEQ ID NO:29)--;
at the end of line 24 insert -- (SEQ ID NO:30)--.

On page 43:

line 14, after "CARRA" insert -- (SEQ ID NO:31)--, and after "RWIPLAK" insert -- (SEQ ID NO:32)--;

00750-492350

PRELIMINARY AMENDMENT

Serial Number: Unknown
Filing Date: Herewith
Title: ARTIFICIAL ANTIBODY POLYPEPTIDES

Page 3
Dkt: 109.034US3

line 15, after "CWRRA" insert -- (SEQ ID NO:33)--, and after "RWVGLAW" insert -
- (SEQ ID NO:34)-- ;

line 16, after "CKHRR" insert -- (SEQ ID NO:35)--, and after "FADLWWR" insert --
(SEQ ID NO:36)--;

line 17, after "CRRGR" insert -- (SEQ ID NO:37)--, and after "RGFMWLS" insert --
(SEQ ID NO:38)-- ;

line 18, after "CNWRR" insert -- (SEQ ID NO:39)--, and after "RAYRYRW" insert -
- (SEQ ID NO:40)-- ;

line 19, after "SRLRR" insert -- (SEQ ID NO:41)--, and after "PPWRV" insert --
(SEQ ID NO: 42)--;

line 20, after "ARWTL" insert -- (SEQ ID NO:43)--, and after "RRWW" insert --
(SEQ ID NO:44)--.

line 21, after "GQRTF" insert -- (SEQ ID NO:45)--, and after "RRWWA" insert --
(SEQ ID NO:46)--.

On page 45:

line 4, after "AVTVR" insert -- (SEQ ID NO:47)--, and after "RGDSPAS" insert --
(SEQ ID NO:48)--;

line 6, after "CNWRR" insert -- (SEQ ID NO:49)--, and after "RAYRYRW" insert --
(SEQ ID NO:50)--;

line 7, after "CMWRA" insert -- (SEQ ID NO:51)--, and after "RWGMLRR" insert --
(SEQ ID NO:52)--;

line 8, after "ARMRE" insert -- (SEQ ID NO:53)--, and after "RWLRGRY" insert --
(SEQ ID NO:54)--;

line 9, after "CARRR" insert -- (SEQ ID NO:55)--, and after "RRAGWGW" insert --
(SEQ ID NO:56)--;

line 10, after "CNWRR" insert -- (SEQ ID NO:57)--, and after "RAYRYRW" insert -
- (SEQ ID NO:58)-- ;

line 11, after "RWRER" insert -- (SEQ ID NO:59)--, and after "RHPWTER" insert --
(SEQ ID NO:60)--;

line 12, after "CNWRR" insert -- (SEQ ID NO:61)--, and after "RAYRYRW" insert -

007130-4136360

- (SEQ ID NO:62)-- ;
line 13, after "ERRVP" insert -- (SEQ ID NO:63)--, and after "RLLWQR" insert -- (SEQ ID NO:64)--;
line 14, after "GRGAG" insert -- (SEQ ID NO:65)--, and after "FGSFERR" insert -- (SEQ ID NO:66)--;
line 15, after "CRWTR" insert -- (SEQ ID NO:67)--, and after "RRWFDGA" insert -- (SEQ ID NO:68)--;
line 16, after "CNWRR" insert -- (SEQ ID NO:69)--, and after "RAYRYRW" insert -- (SEQ ID NO:70)--;
line 20, after "AVTVR" insert -- (SEQ ID NO:71)--, and after "GRGDS" insert -- (SEQ ID NO:72)-- ;
line 22, after "GQRTF" insert -- (SEQ ID NO:73)--, and after "RRWWA" insert -- (SEQ ID NO:74)--;
line 23, after "GQRTF" insert -- (SEQ ID NO:75)--, and after "RRWWA" insert -- (SEQ ID NO:76)--;
line 24, after "GQRTF" insert -- (SEQ ID NO:77)--, and after "RRWWA" insert -- (SEQ ID NO:78)--;
line 25, after "LRYRS" insert -- (SEQ ID NO:79)--, and after "GWRWR" insert -- (SEQ ID NO:80)--;
line 26, after "GQRTF" insert -- (SEQ ID NO:81)--, and after "RRWWA" insert -- (SEQ ID NO:82)--;
line 27, after "GQRTF" insert -- (SEQ ID NO:83)--, and after "RRWWA" insert -- (SEQ ID NO:84)--;
line 28, after "LRYRS" insert -- (SEQ ID NO:85)--, and after "GWRWR" insert -- (SEQ ID NO:86)--;
line 29, after "LRYRS" insert -- (SEQ ID NO:87)--, and after "GWRWR" insert -- (SEQ ID NO:88)--;
line 30, after "GQRTF" insert -- (SEQ ID NO:89)--, and after "RRWWA" insert -- (SEQ ID NO:90)--;
line 31, after "LRYRS" insert -- (SEQ ID NO:91)--, and after "GWRWR" insert --

PRELIMINARY AMENDMENT

Serial Number: Unknown
Filing Date: Herewith
Title: ARTIFICIAL ANTIBODY POLYPEPTIDES

Page 5
Dkt: 109.034US3

(SEQ ID NO:92)--.

On page 52, Table 6:

after "GCAGTTACCGTGCGT" insert -- (SEQ ID NO:93)--, and after
"GGCCGTGGTGACAGCCCAGCGAGC" insert -- (SEQ ID NO:95)--;
after "AlaValThrValArg" insert -- (SEQ ID NO:94)--, and after
"GlyArgGlyAspSerProAlaSer" insert -- (SEQ ID NO:96)-- ;
after "TCGAGGTTGCGGCGG" insert -- (SEQ ID NO:97)--, and after
"CCGCCGTGGAGGGTG" insert -- (SEQ ID NO:99);
after "SerArgLeuArgArg" insert -- (SEQ ID NO:98)--, and after "ProProTrpArgVal"
insert -- (SEQ ID NO:100)-- ;
after "GGTCAGCGAACTTTT" insert -- (SEQ ID NO:101)--, and after
"AGGCGGTGGTGGGCT" insert -- (SEQ ID NO:103);
after "GlyGlnArgThrPhe" insert -- (SEQ ID NO:102)--, and after
"ArgArgTrpTrpAla" insert -- (SEQ ID NO:104)-- ;
after "GCGAGGTGGACGCTT" insert -- (SEQ ID NO:105)--, and after
"AGGCGGTGGTGGTGG" insert -- (SEQ ID NO:107)--;
after "AlaArgTrpThrLeu" insert -- (SEQ ID NO:106)--, and after
"ArgArgTrpTrpTrp" insert -- (SEQ ID NO:108)-- .

On page 54, line 5, after "GKKGK" insert -- (SEQ ID NO:109) --.

In the Claims

Please cancel claims 1-14, 30-35, and 38-39.

In the Drawings

Figure 2 is amended by changing the abbreviation for the amino acid at position 46 from "O" to "Q". Enclosed is a copy of sheet 2 of the drawings showing the amendment to Figure 2 in red ink.

PRELIMINARY AMENDMENT

Serial Number: Unknown
Filing Date: Herewith
Title: ARTIFICIAL ANTIBODY POLYPEPTIDES

Page 6
Dkt: 109.034US3

Remarks

Claims 1-14, 30-35, and 38-39 have been canceled. Claims 15-29, 36 and 37 remain pending.

This Preliminary Amendment is submitted in part to conform the above-referenced application to the requirements of 37 C.F.R. §§ 1.821 through 1.825.

Applicant's Representatives recently discovered that Figure 2 of the Formal Drawings, filed with the application, contains a typographical error. In the first line of the sequence, the "O" at position 46 should be "Q" to represent a glutamine residue. As indicated on page 11, line 25 through page 12, line 2, "O" is not one of the abbreviations set forth whereas "Q" is. Further, amino acid residue number 46 of Figure 5 (SEQ ID NO:111), which corresponds to amino acid residue number 46 of Figure 2, is clearly glutamine. This error will be corrected in the Formal Drawings to be submitted after receipt of a Notice of Allowance.

The Commissioner is hereby authorized to charge any additional fees to Deposit Account No. 19-0743.

When this application is taken up for the first Office Action, Applicant respectfully requests that the Examiner consider the Preliminary Amendment and remarks presented herein.

Respectfully submitted,
SHOHEI KOIDE
By his Representatives,

SCHWEGMAN, LUNDBERG, WOESSNER & KLUTH, P.A.
P.O. Box 2938
Minneapolis, MN 55402
(612) 373-6961

Date 11 August 2000 By Ann S. Viksnins
Ann S. Viksnins
Reg. No. 37,748

Express Mail No.: EL709305975US

Mailing Date: August 11, 2000


I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.

Shawn L. Hise
Printed Name

[Signature]
Signature

[illegible]

FIGURE 3



ARTIFICIAL ANTIBODY POLYPEPTIDES

FIELD OF THE INVENTION

The present invention relates generally to the field of the production and
5 selection of binding and catalytic polypeptides by the methods of molecular biology,
using both combinatorial chemistry and recombinant DNA. The invention
specifically relates to the generation of both nucleic acid and polypeptide libraries
derived therefrom encoding the molecular scaffolding of Fibronectin Type III (Fn3)
modified in one or more of its loop regions. The invention also relates to the
10 “artificial mini-antibodies” or “monobodies,” i.e., the polypeptides comprising an
Fn3 scaffold onto which loop regions capable of binding to a variety of different
molecular structures (such as antibody binding sites) have been grafted.

BACKGROUND OF THE INVENTION

15 Antibody structure

A standard antibody (Ab) is a tetrameric structure consisting of two identical
immunoglobulin (Ig) heavy chains and two identical light chains. The heavy and
light chains of an Ab consist of different domains. Each light chain has one variable
domain (VL) and one constant domain (CL), while each heavy chain has one
20 variable domain (VH) and three or four constant domains (CH) (Alzari et al., 1988).
Each domain, consisting of ~110 amino acid residues, is folded into a characteristic
 β -sandwich structure formed from two β -sheets packed against each other, the
immunoglobulin fold. The VH and VL domains each have three complementarity
determining regions (CDR1-3) that are loops, or turns, connecting β -strands at one
25 end of the domains (Fig. 1: A, C). The variable regions of both the light and heavy
chains generally contribute to antigen specificity, although the contribution of the
individual chains to specificity is not always equal. Antibody molecules have

evolved to bind to a large number of molecules by using six randomized loops (CDRs). However, the size of the antibodies and the complexity of six loops represents a major design hurdle if the end result is to be a relatively small peptide ligand.

5

Antibody substructures

Functional substructures of Abs can be prepared by proteolysis and by recombinant methods. They include the Fab fragment, which comprises the VH-CH1 domains of the heavy chain and the VL-CL1 domains of the light chain joined by a single interchain disulfide bond, and the Fv fragment, which comprises only the VH and VL domains. In some cases, a single VH domain retains significant affinity (Ward et al., 1989). It has also been shown that a certain monomeric κ light chain will specifically bind to its cognate antigen. (L. Masat et al., 1994). Separated light or heavy chains have sometimes been found to retain some antigen-binding activity (Ward et al., 1989). These antibody fragments are not suitable for structural analysis using NMR spectroscopy due to their size, low solubility or low conformational stability.

Another functional substructure is a single chain Fv (scFv), comprised of the variable regions of the immunoglobulin heavy and light chain, covalently connected by a peptide linker (S-z Hu et al., 1996). These small (M_r 25,000) proteins generally retain specificity and affinity for antigen in a single polypeptide and can provide a convenient building block for larger, antigen-specific molecules. Several groups have reported biodistribution studies in xenografted athymic mice using scFv reactive against a variety of tumor antigens, in which specific tumor localization has been observed. However, the short persistence of scFvs in the circulation limits the exposure of tumor cells to the scFvs, placing limits on the level of uptake. As a result, tumor uptake by scFvs in animal studies has generally been only 1-5%ID/g as opposed to intact antibodies that can localize in tumors at 30-40 %ID/g and have reached levels as high as 60-70 %ID/g.

A small protein scaffold called a “minibody” was designed using a part of the Ig VH domain as the template (Pessi et al., 1993). Minibodies with high affinity (dissociation constant (K_d) $\sim 10^{-7}$ M) to interleukin-6 were identified by randomizing loops corresponding to CDR1 and CDR2 of VH and then selecting mutants using the phage display method (Martin et al., 1994). These experiments demonstrated that the essence of the Ab function could be transferred to a smaller system. However, the minibody had inherited the limited solubility of the VH domain (Bianchi et al., 1994).

It has been reported that camels (*Camelus dromedarius*) often lack variable light chain domains when IgG-like material from their serum is analyzed, suggesting that sufficient antibody specificity and affinity can be derived from VH domains (three CDR loops) alone. Davies and Riechmann recently demonstrated that “camelized” VH domains with high affinity ($K_d \sim 10^{-7}$ M) and high specificity can be generated by randomizing only the CDR3. To improve the solubility and suppress nonspecific binding, three mutations were introduced to the framework region (Davies & Riechmann, 1995). It has not been definitively shown, however, that camelization can be used, in general, to improve the solubility and stability of VHs.

An alternative to the “minibody” is the “diabody.” Diabodies are small bivalent and bispecific antibody fragments, i.e., they have two antigen-binding sites. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) on the same polypeptide chain (V_H - V_L). Diabodies are similar in size to an Fab fragment. By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. These dimeric antibody fragments, or “diabodies,” are bivalent and bispecific. P. Holliger et al., PNAS 90:6444-6448 (1993).

Since the development of the monoclonal antibody technology, a large number of 3D structures of Ab fragments in the complexed and/or free states have

been solved by X-ray crystallography (Webster et al., 1994; Wilson & Stanfield, 1994). Analysis of Ab structures has revealed that five out of the six CDRs have limited numbers of peptide backbone conformations, thereby permitting one to predict the backbone conformation of CDRs using the so-called canonical structures (Lesk & Tramontano, 1992; Rees et al., 1994). The analysis also has revealed that the CDR3 of the VH domain (VH-CDR3) usually has the largest contact surface and that its conformation is too diverse for canonical structures to be defined; VH-CDR3 is also known to have a large variation in length (Wu et al., 1993). Therefore, the structures of crucial regions of the Ab-antigen interface still need to be experimentally determined.

Comparison of crystal structures between the free and complexed states has revealed several types of conformational rearrangements. They include side-chain rearrangements, segmental movements, large rearrangements of VH-CDR3 and changes in the relative position of the VH and VL domains (Wilson & Stanfield, 1993). In the free state, CDRs, in particular those which undergo large conformational changes upon binding, are expected to be flexible. Since X-ray crystallography is not suited for characterizing flexible parts of molecules, structural studies in the solution state have not been possible to provide dynamic pictures of the conformation of antigen-binding sites.

Mimicking the antibody-binding site

CDR peptides and organic CDR mimetics have been made (Dougall et al., 1994). CDR peptides are short, typically cyclic, peptides which correspond to the amino acid sequences of CDR loops of antibodies. CDR loops are responsible for antibody-antigen interactions. Organic CDR mimetics are peptides corresponding to CDR loops which are attached to a scaffold, e.g., a small organic compound.

CDR peptides and organic CDR mimetics have been shown to retain some binding affinity (Smyth & von Itzstein, 1994). However, as expected, they are too small and too flexible to maintain full affinity and specificity. Mouse CDRs have

been grafted onto the human Ig framework without the loss of affinity (Jones et al., 1986; Riechmann et al., 1988), though this “humanization” does not solve the above-mentioned problems specific to solution studies.

5 Mimicking natural selection processes of Abs

In the immune system, specific Abs are selected and amplified from a large library (affinity maturation). The processes can be reproduced *in vitro* using combinatorial library technologies. The successful display of Ab fragments on the surface of bacteriophage has made it possible to generate and screen a vast number of CDR mutations (McCafferty et al., 1990; Barbas et al., 1991; Winter et al., 1994). An increasing number of Fabs and Fvs (and their derivatives) is produced by this technique, providing a rich source for structural studies. The combinatorial technique can be combined with Ab mimics.

A number of protein domains that could potentially serve as protein scaffolds have been expressed as fusions with phage capsid proteins. Review in Clackson & Wells, Trends Biotechnol. 12:173-184 (1994). Indeed, several of these protein domains have already been used as scaffolds for displaying random peptide sequences, including bovine pancreatic trypsin inhibitor (Roberts et al., PNAS 89:2429-2433 (1992)), human growth hormone (Lowman et al., Biochemistry 30:10832-10838 (1991)), Venturini et al., Protein Peptide Letters 1:70-75 (1994)), and the IgG binding domain of *Streptococcus* (O’Neil et al., Techniques in Protein Chemistry V (Crabb, L., ed.) pp. 517-524, Academic Press, San Diego (1994)). These scaffolds have displayed a single randomized loop or region.

Researchers have used the small 74 amino acid α -amylase inhibitor Tendamistat as a presentation scaffold on the filamentous phage M13 (McConnell and Hoess, 1995). Tendamistat is a β -sheet protein from *Streptomyces tendae*. It has a number of features that make it an attractive scaffold for peptides, including its small size, stability, and the availability of high resolution NMR and X-ray structural data. Tendamistat’s overall topology is similar to that of an

immunoglobulin domain, with two β -sheets connected by a series of loops. In contrast to immunoglobulin domains, the β -sheets of Tendamistat are held together with two rather than one disulfide bond, accounting for the considerable stability of the protein. By analogy with the CDR loops found in immunoglobulins, the loops the Tendamistat may serve a similar function and can be easily randomized by in vitro mutagenesis.

Tendamistat, however, is derived from *Streptomyces tendae*. Thus, while Tendamistat may be antigenic in humans, its small size may reduce or inhibit its antigenicity. Also, Tendamistat's stability is uncertain. Further, the stability that is reported for Tendamistat is attributed to the presence of two disulfide bonds. Disulfide bonds, however, are a significant disadvantage to such molecules in that they can be broken under reducing conditions and must be properly formed in order to have a useful protein structure. Further, the size of the loops in Tendamistat are relatively small, thus limiting the size of the inserts that can be accommodated in the scaffold. Moreover, it is well known that forming correct disulfide bonds in newly synthesized peptides is not straightforward. When a protein is expressed in the cytoplasmic space of *E. coli*, the most common host bacterium for protein overexpression, disulfide bonds are usually not formed, potentially making it difficult to prepare large quantities of engineered molecules.

Thus, there is an on-going need for small, single-chain artificial antibodies for a variety of therapeutic, diagnostic and catalytic applications.

SUMMARY OF THE INVENTION

The invention provides a fibronectin type III (Fn3) polypeptide monobody comprising a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences. One or more of the monobody loop region sequences of the Fn3 polypeptide vary by deletion, insertion or replacement of at least two amino acids from the corresponding loop region sequences in wild-type Fn3. The β -strand domains of the monobody have at least about 50% total amino

acid sequence homology to the corresponding amino acid sequence of wild-type Fn3's β -strand domain sequences. Preferably, one or more of the loop regions of the monobody comprise amino acid residues:

- i) from 15 to 16 inclusive in an AB loop;
- 5 ii) from 22 to 30 inclusive in a BC loop;
- iii) from 39 to 45 inclusive in a CD loop;
- iv) from 51 to 55 inclusive in a DE loop;
- v) from 60 to 66 inclusive in an EF loop; and
- vi) from 76 to 87 inclusive in an FG loop.

10 The invention also provides a nucleic acid molecule encoding a Fn3 polypeptide monobody of the invention, as well as an expression vector comprising said nucleic acid molecule and a host cell comprising said vector.

The invention further provides a method of preparing a Fn3 polypeptide monobody. The method comprises providing a DNA sequence encoding a plurality
15 of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein at least one loop region of said sequence contains a unique restriction enzyme site. The DNA sequence is cleaved at the unique restriction site. Then a preselected DNA segment is inserted into the restriction site. The preselected DNA segment encodes a peptide capable of binding to a specific binding
20 partner (SBP) or a transition state analog compound (TSAC). The insertion of the preselected DNA segment into the DNA sequence yields a DNA molecule which encodes a polypeptide monobody having an insertion. The DNA molecule is then expressed so as to yield the polypeptide monobody.

Also provided is a method of preparing a Fn3 polypeptide monobody, which
25 method comprises providing a replicatable DNA sequence encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein the nucleotide sequence of at least one loop region is known. Polymerase chain reaction (PCR) primers are provided or prepared which are sufficiently complementary to the known loop sequence so as to be hybridizable

under PCR conditions, wherein at least one of the primers contains a modified nucleic acid sequence to be inserted into the DNA sequence. PCR is performed using the replicatable DNA sequence and the primers. The reaction product of the PCR is then expressed so as to yield a polypeptide monobody.

- 5 The invention further provides a method of preparing a Fn3 polypeptide monobody. The method comprises providing a replicatable DNA sequence encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein the nucleotide sequence of at least one loop region is known. Site-directed mutagenesis of at least one loop region is performed
10 so as to create an insertion mutation. The resultant DNA comprising the insertion mutation is then expressed.

- Further provided is a variegated nucleic acid library encoding Fn3 polypeptide monobodies comprising a plurality of nucleic acid species encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop
15 region sequences, wherein one or more of the monobody loop region sequences vary by deletion, insertion or replacement of at least two amino acids from corresponding loop region sequences in wild-type Fn3, and wherein the β -strand domains of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequence of β -strand domain sequences of the wild-type
20 Fn3. The invention also provides a peptide display library derived from the variegated nucleic acid library of the invention. Preferably, the peptide of the peptide display library is displayed on the surface of a bacteriophage, e.g., a M13 bacteriophage or a fd bacteriophage, or virus.

- The invention also provides a method of identifying the amino acid sequence
25 of a polypeptide molecule capable of binding to a specific binding partner (SBP) so as to form a polypeptide:SSP complex, wherein the dissociation constant of the said polypeptide:SBP complex is less than 10^{-6} moles/liter. The method comprises the steps of:

- a) providing a peptide display library of the invention;

- b) contacting the peptide display library of (a) with an immobilized or separable SBP;
- c) separating the peptide:SBP complexes from the free peptides;
- d) causing the replication of the separated peptides of (c) so as to result in a new peptide display library distinguished from that in (a) by having a lowered diversity and by being enriched in displayed peptides capable of binding the SBP;
- e) optionally repeating steps (b), (c), and (d) with the new library of (d); and
- f) determining the nucleic acid sequence of the region encoding the displayed peptide of a species from (d) and hence deducing the peptide sequence capable of binding to the SBP.

The present invention also provides a method of preparing a variegated nucleic acid library encoding Fn3 polypeptide monobodies having a plurality of nucleic acid species each comprising a plurality of loop regions, wherein the species encode a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein one or more of the loop region sequences vary by deletion, insertion or replacement of at least two amino acids from corresponding loop region sequences in wild-type Fn3, and wherein the β -strand domain sequences of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequences of β -strand domain sequences of the wild-type Fn3, comprising the steps of

- a) preparing an Fn3 polypeptide monobody having a predetermined sequence;
- b) contacting the polypeptide with a specific binding partner (SBP) so as to form a polypeptide:SBP complex wherein the dissociation constant of the said polypeptide:SBP complex is less than 10^{-6} moles/liter;

- c) determining the binding structure of the polypeptide:SBP complex by nuclear magnetic resonance spectroscopy or X-ray crystallography; and
- d) preparing the variegated nucleic acid library, wherein the variegation is performed at positions in the nucleic acid sequence which, from the information provided in (c), result in one or more polypeptides with improved binding to the SBP.

Also provided is a method of identifying the amino acid sequence of a polypeptide molecule capable of catalyzing a chemical reaction with a catalyzed rate constant, k_{cat} , and an uncatalyzed rate constant, k_{uncat} , such that the ratio of k_{cat}/k_{uncat} is greater than 10. The method comprises the steps of:

- a) providing a peptide display library of the invention;
- b) contacting the peptide display library of (a) with an immobilized or separable transition state analog compound (TSAC) representing the approximate molecular transition state of the chemical reaction;
- c) separating the peptide:TSAC complexes from the free peptides;
- d) causing the replication of the separated peptides of (c) so as to result in a new peptide display library distinguished from that in (a) by having a lowered diversity and by being enriched in displayed peptides capable of binding the TSAC;
- e) optionally repeating steps (b), (c), and (d) with the new library of (d); and
- f) determining the nucleic acid sequence of the region encoding the displayed peptide of a species from (d) and hence deducing the peptide sequence.

The invention also provides a method of preparing a variegated nucleic acid library encoding Fn3 polypeptide monobodies having a plurality of nucleic acid species each comprising a plurality of loop regions, wherein the species encode a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop

region sequences, wherein one or more of the loop region sequences vary by deletion, insertion or replacement of at least two amino acids from corresponding loop region sequences in wild-type Fn3, and wherein the β -strand domain sequences of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequences of β -strand domain sequences of the wild-type Fn3, comprising the steps of

- a) preparing an Fn3 polypeptide monobody having a predetermined sequence, wherein the polypeptide is capable of catalyzing a chemical reaction with a catalyzed rate constant, k_{cat} , and an uncatalyzed rate constant, k_{uncat} , such that the ratio of k_{cat}/k_{uncat} is greater than 10;
- b) contacting the polypeptide with an immobilized or separable transition state analog compound (TSAC) representing the approximate molecular transition state of the chemical reaction;
- c) determining the binding structure of the polypeptide:TSAC complex by nuclear magnetic resonance spectroscopy or X-ray crystallography; and
- d) preparing the variegated nucleic acid library, wherein the variegation is performed at positions in the nucleic acid sequence which, from the information provided in (c), result in one or more polypeptides with improved binding to or stabilization of the TSAC.

The invention also provides a kit for the performance of any of the methods of the invention. The invention further provides a composition, e.g., a polypeptide, prepared by the use of the kit, or identified by any of the methods of the invention.

- The following abbreviations have been used in describing amino acids, peptides, or proteins: Ala, or A, Alanine; Arg, or R, Arginine; Asn or N, asparagine; Asp, or D, aspartic acid; Cys or C, cysteine; Gln, or Q, glutamine; Glu, or E, glutamic acid; Gly, or G, glycine; His, or H, histidine; Ile, or I, isoleucine; Leu, or L, leucine; Lys, or K, lysine; Met, or M, methionine; Phe, or F, phenylalanine; Pro, or P,

proline; Ser, or S, serine; Thr, or T, threonine; Trp, or W, tryptophan; Tyr, or Y, tyrosine; Val, or V, valine.

The following abbreviations have been used in describing nucleic acids, DNA, or RNA: A, adenosine; T, thymidine; G, guanosine; C, cytosine.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. β -Strand and loop topology (A, B) and MOLSCRIPT representation (C, D; Kraulis, 1991) of the VH domain of anti-lysozyme immunoglobulin D1.3 (A, C; Bhat et al., 1994) and 10th type III domain of human fibronectin (B, D; Main et al., 1992). The locations of complementarity determining regions (CDRs, hypervariable regions) and the integrin-binding Arg-Gly-Asp (RGD) sequence are indicated.

Figure 2. Amino acid sequence and restriction sites of the synthetic Fn3 gene. The residue numbering is according to Main et al. (1992). Restriction enzyme sites designed are shown above the amino acid sequence. β -Strands are denoted by underlines. The N-terminal "mq" sequence has been added for a subsequent cloning into an expression vector. The His-tag (Novagen) fusion protein has an additional sequence, MGSSHHHHHSSGLVPRGSH, preceding the Fn3 sequence shown above.

Figure 3. A, Far UV CD spectra of wild-type Fn3 at 25°C and 90°C. Fn3 (50 μ M) was dissolved in sodium acetate (50 mM, pH 4.6). B, thermal denaturation of Fn3 monitored at 215 nm. Temperature was increased at a rate of 1°C/min.

Figure 4. A, $C\alpha$ trace of the crystal structure of the complex of lysozyme (HEL) and the Fv fragment of the anti-hen egg-white lysozyme (anti-HEL) antibody D1.3 (Bhat et al., 1994). Side chains of the residues 99-102 of VH CDR3, which make contact with HEL, are also shown. B, Contact surface area for each residue of the D1.3 VH-HEL and VH-VL interactions plotted vs. residue number of D1.3 VH. Surface area and secondary structure were determined using the program DSSP (Kabsh and Sander, 1983). C and D, schematic drawings of the β -sheet structure of

the F strand-loop-G strand moieties of D1.3 VH (C) and Fn3 (D). The boxes denote residues in β -strands and ovals those not in strands. The shaded boxes indicate residues of which side chains are significantly buried. The broken lines indicate hydrogen bonds.

5 Figure 5. Designed Fn3 gene showing DNA and amino acid sequences. The amino acid numbering is according to Main et al. (1992). The two loops that were randomized in combinatorial libraries are enclosed in boxes.

 Figure 6. Map of plasmid pAS45. Plasmid pAS45 is the expression vector of His•tag-Fn3.

10 Figure 7. Map of plasmid pAS25. Plasmid pAS25 is the expression vector of Fn3.

 Figure 8. Map of plasmid pAS38. pAS38 is a phagmid vector for the surface display of Fn3.

 Figure 9. (Ubiquitin-1) Characterization of ligand-specific binding of
15 enriched clones using phage enzyme-linked immunosolvent assay (ELISA). Microtiter plate wells were coated with ubiquitin (1 μ g/well; "Ligand (+)) and then blocked with BSA. Phage solution in TBS containing approximately 10^{10} colony forming units (cfu) was added to a well and washed with TBS. Bound phages were detected with anti-phage antibody-POD conjugate (Pharmacia) with Turbo-TMB
20 (Pierce) as a substrate. Absorbance was measured using a Molecular Devices SPECTRAMax 250 microplate spectrophotometer. For a control, wells without the immobilized ligand were used. 2-1 and 2-2 denote enriched clones from Library 2 eluted with free ligand and acid, respectively. 4-1 and 4-2 denote enriched clones from Library 4 eluted with free ligand and acid, respectively.

25 Figure 10. (Ubiquitin-2) Competition phage ELISA of enriched clones. Phage solutions containing approximately 10^{10} cfu were first incubated with free ubiquitin at 4°C for 1 hour prior to the binding to a ligand-coated well. The wells were washed and phages detected as described above.

Figure 11. Competition phage ELISA of ubiquitin-binding monobody 411. Experimental conditions are the same as described above for ubiquitin. The ELISA was performed in the presence of free ubiquitin in the binding solution. The experiments were performed with four different preparations of the same clone.

5 Figure 12. (Fluorescein-1) Phage ELISA of four clones, pLB25.1, pLB25.4, pLB24.1 and pLB24.3. Experimental conditions are the same as ubiquitin-1 above.

Figure 13. (Fluorescein-2) Competition ELISA of the four clones. Experimental conditions are the same as ubiquitin-2 above.

Figure 14. ^1H , ^{15}N -HSQC spectrum of a fluorescence-binding monobody
10 LB25.5. Approximately 20 μM protein was dissolved in 10 mM sodium acetate buffer (pH 5.0) containing 100 mM sodium chloride. The spectrum was collected at 30°C on a Varian Unity INOVA 600 NMR spectrometer.

Figure 15. Characterization of the binding reaction of Ubi4-Fn3 to the target, ubiquitin. (a) Phage ELISA analysis of binding of Ubi4-Fn3 to ubiquitin.
15 The binding of Ubi4-phages to ubiquitin-coated wells was measured. The control experiment was performed with wells containing no ubiquitin.

(b) Competition phage ELISA of Ubi4-Fn3. Ubi4-Fn3-phages were preincubated with soluble ubiquitin at an indicated concentration, followed by the phage ELISA detection in ubiquitin-coated wells.

20 (c) Competition phage ELISA testing the specificity of the Ubi4 clone. The Ubi4 phages were preincubated with 250 $\mu\text{g/ml}$ of soluble proteins, followed by phage ELISA as in (b).

(d) ELISA using free proteins.

Figure 16. Equilibrium unfolding curves for Ubi4-Fn3 (closed symbols) and
25 wild-type Fn3 (open symbols). Squares indicate data measured in TBS (Tris HCl buffer (50 mM, pH 7.5) containing NaCl (150 mM)). Circles indicate data measured in Gly HCl buffer (20 mM, pH 3.3) containing NaCl (300 mM). The curves show the best fit of the transition curve based on the two-state model. Parameters characterizing the transitions are listed in Table 7.

Figure 17. (a) ^1H , ^{15}N -HSQC spectrum of [^{15}N]-Ubi4-K Fn3. (b). Difference ($\delta_{\text{wild-type}} - \delta_{\text{Ubi4}}$) of ^1H (b) and ^{15}N (c) chemical shifts plotted versus residue number. Values for residues 82-84 (shown as filled circles) where Ubi4-K deletions are set to zero. Open circles indicate residues that are mutated in the Ubi4-K protein. The locations of β -strands are indicated with arrows.

DETAILED DESCRIPTION OF THE INVENTION

For the past decade the immune system has been exploited as a rich source of *de novo* catalysts. Catalytic antibodies have been shown to have chemoselectivity, enantioselectivity, large rate accelerations, and even an ability to reroute chemical reactions. In most cases the antibodies have been elicited to transition state analog (TSA) haptens. These TSA haptens are stable, low-molecular weight compounds designed to mimic the structures of the energetically unstable transition state species that briefly (approximate half-life 10^{-13} s) appear along reaction pathways between reactants and products. Anti-TSA antibodies, like natural enzymes, are thought to selectively bind and stabilize transition state, thereby easing the passage of reactants to products. Thus, upon binding, the antibody lowers the energy of the actual transition state and increases the rate of the reaction. These catalysts can be programmed to bind to geometrical and electrostatic features of the transition state so that the reaction route can be controlled by neutralizing unfavorable charges, overcoming entropic barriers, and dictating stereoelectronic features of the reaction. By this means even reactions that are otherwise highly disfavored have been catalyzed (Janda et al. 1997). Further, in many instances catalysts have been made for reactions for which there are no known natural or man-made enzymes.

The success of any combinatorial chemical system in obtaining a particular function depends on the size of the library and the ability to access its members. Most often the antibodies that are made in an animal against a hapten that mimics the transition state of a reaction are first screened for binding to the hapten and then

screened again for catalytic activity. An improved method allows for the direct selection for catalysis from antibody libraries in phage, thereby linking chemistry and replication.

5 A library of antibody fragments can be created on the surface of filamentous phage viruses by adding randomized antibody genes to the gene that encodes the phage's coat protein. Each phage then expresses and displays multiple copies of a single antibody fragment on its surface. Because each phage possesses both the surface-displayed antibody fragment and the DNA that encodes that fragment, and antibody fragment that binds to a target can be identified by amplifying the
10 associated DNA.

Immunochemists use as antigens materials that have as little chemical reactivity as possible. It is almost always the case that one wishes the ultimate antibody to interact with native structures. In reactive immunization the concept is just the opposite. One immunizes with compounds that are highly reactive so that
15 upon binding to the antibody molecule during the induction process, a chemical reaction ensues. Later this same chemical reaction becomes part of the mechanism of the catalytic event. In a certain sense one is immunizing with a chemical reaction rather than a substance *per se*. Reactive immunogens can be considered as analogous to the mechanism-based inhibitors that enzymologists use except that
20 they are used in the inverse way in that, instead of inhibiting a mechanism, they induce a mechanism.

Man-made catalytic antibodies have considerable commercial potential in many different applications. Catalytic antibody-based products have been used successfully in prototype experiments in therapeutic applications, such as prodrug
25 activation and cocaine inactivation, and in nontherapeutic applications, such as biosensors and organic synthesis.

Catalytic antibodies are theoretically more attractive than noncatalytic antibodies as therapeutic agents because, being catalytic, they may be used in lower doses, and also because their effects are unusually irreversible (for example, peptide

bond cleavage rather than binding). In therapy, purified catalytic antibodies could be directly administered to a patient, or alternatively the patient's own catalytic antibody response could be elicited by immunization with an appropriate hapten. Catalytic antibodies also could be used as clinical diagnostic tools or as
5 regioselective or stereoselective catalysts in the synthesis of fine chemicals.

I. Mutation of Fn3 loops and grafting of Ab loops onto Fn3

An ideal scaffold for CDR grafting is highly soluble and stable. It is small enough for structural analysis, yet large enough to accommodate multiple CDRs so
10 as to achieve tight binding and/or high specificity.

A novel strategy to generate an artificial Ab system on the framework of an existing non-Ab protein was developed. An advantage of this approach over the minimization of an Ab scaffold is that one can avoid inheriting the undesired properties of Abs. Fibronectin type III domain (Fn3) was used as the scaffold.
15 Fibronectin is a large protein which plays essential roles in the formation of extracellular matrix and cell-cell interactions; it consists of many repeats of three types (I, II and III) of small domains (Baron et al., 1991). Fn3 itself is the paradigm of a large subfamily (Fn3 family or s-type Ig family) of the immunoglobulin superfamily (IgSF). The Fn3 family includes cell adhesion molecules, cell surface
20 hormone and cytokine receptors, chaperonins, and carbohydrate-binding domains (for reviews, see Bork & Doolittle, 1992; Jones, 1993; Bork et al., 1994; Campbell & Spitzfaden, 1994; Harpez & Chothia, 1994).

Recently, crystallographic studies revealed that the structure of the DNA binding domains of the transcription factor NF- κ B is also closely related to the Fn3
25 fold (Ghosh et al., 1995; Müller et al., 1995). These proteins are all involved in specific molecular recognition, and in most cases ligand-binding sites are formed by surface loops, suggesting that the Fn3 scaffold is an excellent framework for building specific binding proteins. The 3D structure of Fn3 has been determined by NMR (Main et al., 1992) and by X-ray crystallography (Leahy et al., 1992;

Dickinson et al., 1994). The structure is best described as a β -sandwich similar to that of Ab VH domain except that Fn3 has seven β -strands instead of nine (Fig. 1). There are three loops on each end of Fn3; the positions of the BC, DE and FG loops approximately correspond to those of CDR1, 2 and 3 of the VH domain,

5 respectively (Fig. 1 C, D).

Fn3 is small (~ 95 residues), monomeric, soluble and stable. It is one of few members of IgSF that do not have disulfide bonds; VH has an interstrand disulfide bond (Fig. 1 A) and has marginal stability under reducing conditions. Fn3 has been expressed in *E. coli* (Aukhil et al., 1993). In addition, 17 Fn3 domains are present
10 just in human fibronectin, providing important information on conserved residues which are often important for the stability and folding (for sequence alignment, see Main et al., 1992 and Dickinson et al., 1994). From sequence analysis, large variations are seen in the BC and FG loops, suggesting that the loops are not crucial to stability. NMR studies have revealed that the FG loop is highly flexible; the
15 flexibility has been implicated for the specific binding of the 10th Fn3 to $\alpha_5\beta_1$ integrin through the Arg-Gly-Asp (RGD) motif. In the crystal structure of human growth hormone-receptor complex (de Vos et al., 1992), the second Fn3 domain of the receptor interacts with hormone via the FG and BC loops, suggesting it is feasible to build a binding site using the two loops.

20 The tenth type III module of fibronectin has a fold similar to that of immunoglobulin domains, with seven β strands forming two antiparallel β sheets, which pack against each other (Main et al., 1992). The structure of the type II module consists of seven β strands, which form a sandwich of two antiparallel β sheets, one containing three strands (ABE) and the other four strands (C'CFG)
25 (Williams et al., 1988). The triple-stranded β sheet consists of residues Glu-9-Thr-14 (A), Ser-17-Asp-23 (B), and Thr-56-Ser-60 (E). The majority of the conserved residues contribute to the hydrophobic core, with the invariant hydrophobic residues Trp-22 and Try-68 lying toward the N-terminal and C-terminal ends of the core, respectively. The β strands are much less flexible and appear to provide a rigid

framework upon which functional, flexible loops are built. The topology is similar to that of immunoglobulin C domains.

Gene construction and mutagenesis

5 A synthetic gene for tenth Fn3 of human fibronectin (Fig. 2) was designed which includes convenient restriction sites for ease of mutagenesis and uses specific codons for high-level protein expression (Gribskov et al., 1984).

10 The gene was assembled as follows: (1) the gene sequence was divided into five parts with boundaries at designed restriction sites (Fig.2); (2) for each part, a pair of oligonucleotides that code opposite strands and have complementary overlaps of ~ 15 bases was synthesized; (3) the two oligonucleotides were annealed and single strand regions were filled in using the Klenow fragment of DNA polymerase; (4) the double-stranded oligonucleotide was cloned into the pET3a vector (Novagen) using restriction enzyme sites at the termini of the fragment and
15 its sequence was confirmed by an Applied Biosystems DNA sequencer using the dideoxy termination protocol provided by the manufacturer; (5) steps 2-4 were repeated to obtain the whole gene (plasmid pAS25) (Fig. 7).

20 Although the present method takes more time to assemble a gene than the one-step polymerase chain reaction (PCR) method (Sandhu et al., 1992), no mutations occurred in the gene. Mutations would likely have been introduced by the low fidelity replication by Taq polymerase and would have required time-consuming gene editing. The gene was also cloned into the pET15b (Novagen) vector (pEW1). Both vectors expressed the Fn3 gene under the control of bacteriophage T7 promoter (Studler et al. 1990); pAS25 expressed the 96-residue Fn3 protein only, while pEW1
25 expressed Fn3 as a fusion protein with poly-histidine peptide (His•tag). Recombinant DNA manipulations were performed according to Molecular Cloning (Sambrook et al., 1989), unless otherwise stated.

 Mutations were introduced to the Fn3 gene using either cassette mutagenesis or oligonucleotide site-directed mutagenesis techniques (Deng & Nickoloff, 1992).

Cassette mutagenesis was performed using the same protocol for gene construction described above; double-stranded DNA fragment coding a new sequence was cloned into an expression vector (pAS25 and/or pEW1). Many mutations can be made by combining a newly synthesized strand (coding mutations) and an oligonucleotide used for the gene synthesis. The resulting genes were sequenced to confirm that the designed mutations and no other mutations were introduced by mutagenesis reactions.

Design and synthesis of Fn3 mutants with antibody CDRs

Two candidate loops (FG and BC) were identified for grafting. Antibodies with known crystal structures were examined in order to identify candidates for the sources of loops to be grafted onto Fn3. Anti-hen egg lysozyme (HEL) antibody D1.3 (Bhat et al., 1994) was chosen as the source of a CDR loop. The reasons for this choice were: (1) high resolution crystal structures of the free and complexed states are available (Fig. 4 A; Bhat et al., 1994), (2) thermodynamics data for the binding reaction are available (Tello et al., 1993), (3) D1.3 has been used as a paradigm for Ab structural analysis and Ab engineering (Verhoeyen et al., 1988; McCafferty et al., 1990) (4) site-directed mutagenesis experiments have shown that CDR3 of the heavy chain (VH-CDR3) makes a larger contribution to the affinity than the other CDRs (Hawkins et al., 1993), and (5) a binding assay can be easily performed. The objective for this trial was to graft VH-CDR3 of D1.3 onto the Fn3 scaffold without significant loss of stability.

An analysis of the D1.3 structure (Fig. 4) revealed that only residues 99-102 ("RDYR") make direct contact with hen egg-white lysozyme (HEL) (Fig. 4 B), although VH-CDR3 is defined as longer (Bhat et al., 1994). It should be noted that the C-terminal half of VH-CDR3 (residues 101-104) made significant contact with the VL domain (Fig. 4 B). It has also become clear that D1.3 VH-CDR3 (Fig. 4 C) has a shorter turn between the strands F and G than the FG loop of Fn3 (Fig. 4 D). Therefore, mutant sequences were designed by using the RDYR (99-102) of D1.3 as

the core and made different boundaries and loop lengths (Table 1). Shorter loops may mimic the D1.3 CDR3 conformation better, thereby yielding higher affinity, but they may also significantly reduce stability by removing wild-type interactions of Fn3.

5

Table 1. Amino acid sequences of D1.3 VH CDR3, VH8 CDR3 and Fn3 FG loop and list of planned mutants.

		96	100	105
		•	•	•
	D1.3	<u>A R E R D Y R L D Y W G Q G</u>		
10	VH8	<u>A R G A V V S Y Y A M D Y W G Q G</u>		
		75	80	85
		•	•	•
	Fn3	<u>Y A V T G R G D S P A S S K P I</u>		
	Mutant	Sequence		
	D1.3-1	Y A E R D Y R L D Y ---- P I		
	D1.3-2	Y A V R D Y R L D Y ---- P I		
15	D1.3-3	Y A V R D Y R L D Y A S S K P I		
	D1.3-4	Y A V R D Y R L D Y --- K P I		
	D1.3-5	Y A V R D Y R ----- S K P I		
	D1.3-6	Y A V T R D Y R L -- S S K P I		
	D1.3-7	Y A V T E R D Y R L - S S K P I		
20	VH8-1	Y A V A V V S Y Y A M D Y - P I		
	VH8-2	Y A V T A V V S Y Y A S S K P I		

Underlines indicate residues in β -strands. Bold characters indicate replaced residues.

In addition, an anti-HEL single VH domain termed VH8 (Ward et al., 1989) was chosen as a template. VH8 was selected by library screening and, in spite of the lack of the VL domain, VH8 has an affinity for HEL of 27 nM, probably due to its longer VH-CDR3 (Table 1). Therefore, its VH-CDR3 was grafted onto Fn3.

- 5 Longer loops may be advantageous on the Fn3 framework because they may provide higher affinity and also are close to the loop length of wild-type Fn3. The 3D structure of VH8 was not known and thus the VH8 CDR3 sequence was aligned with that of D1.3 VH-CDR3; two loops were designed (Table 1).

10 **Mutant construction and production**

Site-directed mutagenesis experiments were performed to obtain designed sequences. Two mutant Fn3s, D1.3-1 and D1.3-4 (Table 1) were obtained and both were expressed as soluble His•tag fusion proteins. D1.3-4 was purified and the His•tag portion was removed by thrombin cleavage. D1.3-4 is soluble up to at least
15 1 mM at pH 7.2. No aggregation of the protein has been observed during sample preparation and NMR data acquisition.

Protein expression and purification

- E. coli* BL21 (DE3) (Novagen) were transformed with an expression vector
20 (pAS25, pEW1 and their derivatives) containing a gene for the wild-type or a mutant. Cells were grown in M9 minimal medium and M9 medium supplemented with Bactotrypton (Difco) containing ampicillin (200 µg/ml). For isotopic labeling, ¹⁵N NH₄Cl and/or ¹³C glucose replaced unlabeled components. 500 ml medium in a 2 liter baffie flask were inoculated with 10 ml of overnight culture and agitated at
25 37°C. Isopropylthio-β-galactoside (IPTG) was added at a final concentration of 1 mM to initiate protein expression when OD (600 nm) reaches one. The cells were harvested by centrifugation 3 hours after the addition of IPTG and kept frozen at -70°C until used.

Fn3 without His•tag was purified as follows. Cells were suspended in 5 ml/(g cell) of Tris (50 mM, pH 7.6) containing ethylenediaminetetraacetic acid (EDTA; 1 mM) and phenylmethylsulfonyl fluoride (1 mM). HEL was added to a final concentration of 0.5 mg/ml. After incubating the solution for 30 minutes at 37°C, it was sonicated three times for 30 seconds on ice. Cell debris was removed by centrifugation. Ammonium sulfate was added to the solution and precipitate recovered by centrifugation. The pellet was dissolved in 5-10 ml sodium acetate (50 mM, pH 4.6) and insoluble material was removed by centrifugation. The solution was applied to a Sephacryl S100HR column (Pharmacia) equilibrated in the sodium acetate buffer. Fractions containing Fn3 then was applied to a ResourceS column (Pharmacia) equilibrated in sodium acetate (50 mM, pH 4.6) and eluted with a linear gradient of sodium chloride (0-0.5 M). The protocol can be adjusted to purify mutant proteins with different surface charge properties.

Fn3 with His•tag was purified as follows. The soluble fraction was prepared as described above, except that sodium phosphate buffer (50 mM, pH 7.6) containing sodium chloride (100 mM) replaced the Tris buffer. The solution was applied to a Hi-Trap chelating column (Pharmacia) preloaded with nickel and equilibrated in the phosphate buffer. After washing the column with the buffer, His•tag-Fn3 was eluted in the phosphate buffer containing 50 mM EDTA. Fractions containing His•tag-Fn3 were pooled and applied to a Sephacryl S100-HR column, yielding highly pure protein. The His•tag portion was cleaved off by treating the fusion protein with thrombin using the protocol supplied by Novagen. Fn3 was separated from the His•tag peptide and thrombin by a ResourceS column using the protocol above.

The wild-type and two mutant proteins so far examined are expressed as soluble proteins. In the case that a mutant is expressed as inclusion bodies (insoluble aggregate), it is first examined if it can be expressed as a soluble protein at lower temperature (e.g., 25-30°C). If this is not possible, the inclusion bodies are collected by low-speed centrifugation following cell lysis as described above. The

pellet is washed with buffer, sonicated and centrifuged. The inclusion bodies are solubilized in phosphate buffer (50 mM, pH 7.6) containing guanidinium chloride (GdnCl, 6 M) and will be loaded on a Hi-Trap chelating column. The protein is eluted with the buffer containing GdnCl and 50 mM EDTA.

5

Conformation of mutant Fn3, D1.3-4

The ¹H NMR spectra of His•tag D1.3-4 fusion protein closely resembled that of the wild-type, suggesting the mutant is folded in a similar conformation to that of the wild-type. The spectrum of D1.3-4 after the removal of the His•tag peptide
10 showed a large spectral dispersion. A large dispersion of amide protons (7-9.5 ppm) and a large number of downfield (5.0-6.5 ppm) C^α protons are characteristic of a β-sheet protein (Wüthrich, 1986).

The 2D NOESY spectrum of D1.3-4 provided further evidence for a preserved conformation. The region in the spectrum showed interactions between
15 upfield methyl protons (< 0.5 ppm) and methyl-methylene protons. The Val72 γ methyl resonances were well separated in the wild-type spectrum (-0.07 and 0.37 ppm; (Baron et al., 1992)). Resonances corresponding to the two methyl protons are present in the D1.3-4 spectrum (-0.07 and 0.44 ppm). The cross peak between these two resonances and other conserved cross peaks indicate that the two
20 resonances in the D1.3-4 spectrum are highly likely those of Val72 and that other methyl protons are in nearly identical environment to that of wild-type Fn3. Minor differences between the two spectra are presumably due to small structural perturbation due to the mutations. Val72 is on the F strand, where it forms a part of the central hydrophobic core of Fn3 (Main et al., 1992). It is only four residues
25 away from the mutated residues of the FG loop (Table 1). The results are remarkable because, despite there being 7 mutations and 3 deletions in the loop (more than 10% of total residues; Fig. 12, Table 2), D1.3-4 retains a 3D structure virtually identical to that of the wild-type (except for the mutated loop). Therefore, the results provide strong support that the FG loop is not significantly contributing

to the folding and stability of the Fn3 molecule and thus that the FG loop can be mutated extensively.

Table 2. Sequences of oligonucleotides

5		
	Name	Sequence
	FN1F	CGGGATCC <u>CATATG</u> CAGGTTTCTGATGTTCCGCGTGACCTGGAAGTTGTTGCTGCGACC
	FN1R	TA <u>ACTG</u> CAGGAGCATCCCAGCTGATCAGCAGGCTAGTCGGGGTCGCAGCAACAAC
	FN2F	CTC <u>CTGCAG</u> TTACCGTGC GTTATTACCGTATCACGTACGGTGAAACCGGTG
10	FN2R	GTGAATTCCTGAACCGGGGAGTTACCACCGGTTTCACCG
	FN3F	AGGAATTCACCTGTACCTGGTTCCAAGTCTACTGCTACCATCAGCGG
	FN3R	GTATAGTCGACACCCGGTTTCAGGCCGCTGATGGTAGC
	FN4F	CGGGTGT <u>CGACT</u> TATACCATCACTGTATACGCT
	FN4R	CGGGATCCGAGCTCGCTGGGCTGTCACCACGGCCAGTAACAGCGTATACAGTGAT
15	FN5F	CAGCGAGCTCCAAGCCAATCTCGATTAACTACCGT
	FN5R	CGGGATCCCTCGAGTTACTAGGTACGGTAGTTAATCGA
	FN5R'	CGGGATCCACGCGTGCCACCGGTACGGTAGTTAATCGA
	gene3F	CGGGATCCACGCGTCCATTCGTTTGTGAATATCAAGGCCAATCG
	gene3R	CCGGAAGCTTTAAGACTCCTTATTACGCAGTATGTTAGC
20	38TAABgII	CTGTTACTGGCCGTGAGATCTAACCAGCGAGCTCCA
	BC3	GATCAGCTGGGATGCTCCTNNKNNKNNKNNKNNKTATTACCGTATCACGTA
	FG2	TGTATACGCTGTTACTGGCNNKNNKNNKNNKNNKNNKNNKTCCAAGCCAATCTCGAT
	FG3	CTGTATACGCTGTTACTGGCNNKNNKNNKNNKCCAGCGAGCTCCAAG
	FG4	CATCACTGTATACGCTGTTACTNNKNNKNNKNNKNNKTCCAAGCCAATCTC

25

Restriction enzyme sites are underlined. N and K denote an equimolar mixture of A, T, G and C and that of G and T, respectively.

30 Structure and stability measurements

Structures of Abs were analyzed using quantitative methods (e.g., DSSP (Kabsch & Sander, 1983) and PDBfit (D. McRee, The Scripps Research Institute)) as well as computer graphics (e.g., Quanta (Molecular Simulations) and What if (G. Vriend, European Molecular Biology Laboratory)) to superimpose the strand-loop-strand structures of Abs and Fn3.

35

006374-09100
00T30"4T92960

The stability of FnAbs was determined by measuring temperature- and chemical denaturant-induced unfolding reactions (Pace et al., 1989). The temperature-induced unfolding reaction was measured using a circular dichroism (CD) polarimeter. Ellipticity at 222 and 215 nm was recorded as the sample temperature was slowly raised. Sample concentrations between 10 and 50 μ M were used. After the unfolding baseline was established, the temperature was lowered to examine the reversibility of the unfolding reaction. Free energy of unfolding was determined by fitting data to the equation for the two-state transition (Becktel & Schellman, 1987; Pace et al., 1989). Nonlinear least-squares fitting was performed using the program Igor (WaveMetrics) on a Macintosh computer.

The structure and stability of two selected mutant Fn3s were studied; the first mutant was D1.3-4 (Table 2) and the second was a mutant called AS40 which contains four mutations in the BC loop ($A^{26}V^{27}T^{28}V^{29} \rightarrow TQRQ$). AS40 was randomly chosen from the BC loop library described above. Both mutants were expressed as soluble proteins in *E. coli* and were concentrated at least to 1 mM, permitting NMR studies.

The mid-point of the thermal denaturation for both mutants was approximately 69°C, as compared to approximately 79°C for the wild-type protein. The results indicated that the extensive mutations at the two surface loops did not drastically decrease the stability of Fn3, and thus demonstrated the feasibility of introducing a large number of mutations in both loops.

Stability was also determined by guanidinium chloride (GdnCl)- and urea-induced unfolding reactions. Preliminary unfolding curves were recorded using a fluorometer equipped with a motor-driven syringe; GdnCl or urea were added continuously to the protein solution in the cuvette. Based on the preliminary unfolding curves, separate samples containing varying concentration of a denaturant were prepared and fluorescence (excitation at 290 nm, emission at 300-400 nm) or CD (ellipticity at 222 and 215 nm) were measured after the samples were equilibrated at the measurement temperature for at least one hour. The curve was

09637614-081100
fitted by the least-squares method to the equation for the two-state model (Santoro & Bolen, 1988; Koide et al., 1993). The change in protein concentration was compensated if required.

Once the reversibility of the thermal unfolding reaction is established, the
5 unfolding reaction is measured by a Microcal MC-2 differential scanning
calorimeter (DSC). The cell (~ 1.3 ml) will be filled with FnAb solution (0.1 -
1 mM) and $\Delta C_p (= \Delta H/\Delta T)$ will be recorded as the temperature is slowly raised. T_m
(the midpoint of unfolding), ΔH of unfolding and ΔG of unfolding is determined by
fitting the transition curve (Privalov & Potekhin, 1986) with the Origin software
10 provided by Microcal.

Thermal unfolding

A temperature-induced unfolding experiment on Fn3 was performed using
circular dichroism (CD) spectroscopy to monitor changes in secondary structure.
15 The CD spectrum of the native Fn3 shows a weak signal near 222 nm (Fig. 3A),
consistent with the predominantly β -structure of Fn3 (Perczel et al., 1992). A
cooperative unfolding transition is observed at 80-90°C, clearly indicating high
stability of Fn3 (Fig. 3B). The free energy of unfolding could not be determined due
to the lack of a post-transition baseline. The result is consistent with the high
20 stability of the first Fn3 domain of human fibronectin (Litvinovich et al., 1992), thus
indicating that Fn3 domains are in general highly stable.

Binding assays

Binding reaction of FnAbs were characterized quantitatively using an
25 isothermal titration calorimeter (ITC) and fluorescence spectroscopy.

The enthalpy change (ΔH) of binding were measured using a Microcal
Omega ITC (Wiseman et al., 1989). The sample cell (~ 1.3 ml) was filled with
FnAbs solution ($\leq 100 \mu\text{M}$, changed according to K_d), and the reference cell filled
with distilled water; the system was equilibrated at a given temperature until a stable

baseline is obtained; 5-20 μ l of ligand solution (≤ 2 mM) was injected by a motor-driven syringe within a short duration (20 sec) followed by an equilibration delay (4 minutes); the injection was repeated and heat generation/absorption for each injection was measured. From the change in the observed heat change as a function of ligand concentration, ΔH and K_d was determined (Wiseman et al., 1989). ΔG and ΔS of the binding reaction was deduced from the two directly measured parameters. Deviation from the theoretical curve was examined to assess nonspecific (multiple-site) binding. Experiments were also performed by placing a ligand in the cell and titrating with an FnAb. It should be emphasized that only ITC gives direct measurement of ΔH , thereby making it possible to evaluate enthalpic and entropic contributions to the binding energy. ITC was successfully used to monitor the binding reaction of the D1.3 Ab (Tello et al., 1993; Bhat et al., 1994).

Intrinsic fluorescence is monitored to measure binding reactions with K_d in the sub- μ M range where the determination of K_d by ITC is difficult. Trp fluorescence (excitation at ~ 290 nm, emission at 300-350 nm) and Tyr fluorescence (excitation at ~ 260 nm, emission at ~ 303 nm) is monitored as the Fn3-mutant solution (≤ 10 μ M) is titrated with ligand solution (≤ 100 μ M). K_d of the reaction is determined by the nonlinear least-squares fitting of the bimolecular binding equation. Presence of secondary binding sites is examined using Scatchard analysis. In all binding assays, control experiments are performed using wild-type Fn3 (or unrelated FnAbs) in place of FnAbs of interest.

II. Production of Fn3 mutants with high affinity and specificity FnAbs

Library screening was carried out in order to select FnAbs which bind to specific ligands. This is complementary to the modeling approach described above. The advantage of combinatorial screening is that one can easily produce and screen a large number of variants ($\geq 10^8$), which is not feasible with specific mutagenesis ("rational design") approaches. The phage display technique (Smith, 1985; O'Neil & Hoess, 1995) was used to effect the screening processes. Fn3 was fused to a

phage coat protein (pIII) and displayed on the surface of filamentous phages. These phages harbor a single-stranded DNA genome that contains the gene coding the Fn3 fusion protein. The amino acid sequence of defined regions of Fn3 were randomized using a degenerate nucleotide sequence, thereby constructing a library.

- 5 Phages displaying Fn3 mutants with desired binding capabilities were selected *in vitro*, recovered and amplified. The amino acid sequence of a selected clone can be identified readily by sequencing the Fn3 gene of the selected phage. The protocols of Smith (Smith & Scott, 1993) were followed with minor modifications.

- 10 The objective was to produce FnAbs which have high affinity to small protein ligands. HEL and the B1 domain of staphylococcal protein G (hereafter referred to as protein G) were used as ligands. Protein G is small (56 amino acids) and highly stable (Minor & Kim, 1994; Smith et al., 1994). Its structure was determined by NMR spectroscopy (Gronenborn et al., 1991) to be a helix packed against a four-strand β -sheet. The resulting FnAb-protein G complexes (~ 150
15 residues) is one of the smallest protein-protein complexes produced to date, well within the range of direct NMR methods. The small size, the high stability and solubility of both components and the ability to label each with stable isotopes (^{13}C and ^{15}N ; see below for protein G) make the complexes an ideal model system for NMR studies on protein-protein interactions.

- 20 The successful loop replacement of Fn3 (the mutant D1.3-4) demonstrate that at least ten residues can be mutated without the loss of the global fold. Based on this, a library was first constructed in which only residues in the FG loop are randomized. After results of loop replacement experiments on the BC loop were obtained, mutation sites were extended that include the BC loop and other sites.

25

Construction of Fn3 phage display system

An M13 phage-based expression vector pASM1 has been constructed as follows: an oligonucleotide coding the signal peptide of OmpT was cloned at the 5' end of the Fn3 gene; a gene fragment coding the C-terminal domain of M13 pIII was

prepared from the wild-type gene III gene of M13 mp18 using PCR (Corey et al., 1993) and the fragment was inserted at the 3' end of the OmpT-Fn3 gene; a spacer sequence has been inserted between Fn3 and pIII. The resultant fragment (OmpT-Fn3-pIII) was cloned in the multiple cloning site of M13 mp18, where the fusion gene is under the control of the lac promoter. This system will produce the Fn3-pIII fusion protein as well as the wild-type pIII protein. The co-expression of wild-type pIII is expected to reduce the number of fusion pIII protein, thereby increasing the phage infectivity (Corey et al., 1993) (five copies of pIII are present on a phage particle). In addition, a smaller number of fusion pIII protein may be advantageous in selecting tight binding proteins, because the chelating effect due to multiple binding sites should be smaller than that with all five copies of fusion pIII (Bass et al., 1990). This system has successfully displayed the serine protease trypsin (Corey et al., 1993). Phages were produced and purified using *E. coli* K91kan (Smith & Scott, 1993) according to a standard method (Sambrook et al., 1989) except that phage particles were purified by a second polyethylene glycol precipitation and acid precipitation.

Successful display of Fn3 on fusion phages has been confirmed by ELISA using an Ab against fibronectin (Sigma), clearly indicating that it is feasible to construct libraries using this system.

An alternative system using the fUSE5 (Parmley & Smith, 1988) may also be used. The Fn3 gene is inserted to fUSE5 using the SfiI restriction sites introduced at the 5'- and 3'- ends of the Fn3 gene PCR. This system displays only the fusion pIII protein (up to five copies) on the surface of a phage. Phages are produced and purified as described (Smith & Scott, 1993). This system has been used to display many proteins and is robust. The advantage of fUSE5 is its low toxicity. This is due to the low copy number of the replication form (RF) in the host, which in turn makes it difficult to prepare a sufficient amount of RF for library construction (Smith & Scott, 1993).

Construction of libraries

The first library was constructed of the Fn3 domain displayed on the surface of MB phage in which seven residues (77-83) in the FG loop (Fig. 4D) were randomized. Randomization will be achieved by the use of an oligonucleotide containing degenerated nucleotide sequence. A double-stranded nucleotide was prepared by the same protocol as for gene synthesis (see above) except that one strand had an (NNK)₆(NNG) sequence at the mutation sites, where N corresponds to an equimolar mixture of A, T, G and C and K corresponds to an equimolar mixture of G and T. The (NNG) codon at residue 83 was required to conserve the SacI restriction site (Fig. 2). The (NNK) codon codes all of the 20 amino acids, while the NNG codon codes 14. Therefore, this library contained ~ 10⁹ independent sequences. The library was constructed by ligating the double-stranded nucleotide into the wild-type phage vector, pASM1, and the transfecting *E. coli* XL1 blue (Stratagene) using electroporation. XL1 blue has the lacI^q phenotype and thus suppresses the expression of the Fn3-pIII fusion protein in the absence of lac inducers. The initial library was propagated in this way, to avoid selection against toxic Fn3-pIII clones. Phages displaying the randomized Fn3-pIII fusion protein were prepared by propagating phages with K91kan as the host. K91kan does not suppress the production of the fusion protein, because it does not have lacI^q. Another library was also generated in which the BC loop (residues 26-30) was randomized.

Selection of displayed FnAbs

Screening of Fn3 phage libraries was performed using the biopanning protocol (Smith & Scott, 1993); a ligand is biotinylated and the strong biotin-streptavidin interaction was used to immobilize the ligand on a streptavidin-coated dish. Experiments were performed at room temperature (~ 22°C). For the initial recovery of phages from a library, 10 µg of a biotinylated ligand were immobilized on a streptavidin-coated polystyrene dish (35 mm, Falcon 1008) and then a phage

007750-4794650
tetramethylbenzidine, Pierce). The high affinity of protein G to immunoglobulins present a special problem; Abs cannot be used in detection. Therefore, to detect anti-protein G FnAbs, fusion phages are immobilized in wells and the binding is then measured using biotinylated protein G followed by the detection using streptavidin-peroxidase conjugate.

Production of soluble FnAbs

After preliminary characterization of mutant Fn3s using phage ELISA, mutant genes are subcloned into the expression vector pEW1. Mutant proteins are produced as His•tag fusion proteins and purified, and their conformation, stability and ligand affinity are characterized.

Thus, Fn3 is the fourth example of a monomeric immunoglobulin-like scaffold that can be used for engineering binding proteins. Successful selection of novel binding proteins have also been based on minibody, tendamistat and “camelized” immunoglobulin VH domain scaffolds (Martin et al., 1994; Davies & Riechmann, 1995; McConnell & Hoess, 1995). The Fn3 scaffold has advantages over these systems. Bianchi et al. reported that the stability of a minibody was 2.5 kcal/mol, significantly lower than that of Ubi4-K. No detailed structural characterization of minibodies has been reported to date. Tendamistat and the VH domain contain disulfide bonds, and thus preparation of correctly folded proteins may be difficult. Davies and Riechmann reported that the yields of their camelized VH domains were less than 1 mg per liter culture (Davies & Riechmann, 1996).

Thus, the Fn3 framework can be used as a scaffold for molecular recognition. Its small size, stability and well-characterized structure make Fn3 an attractive system. In light of the ubiquitous presence of Fn3 in a wide variety of natural proteins involved in ligand binding, one can engineer Fn3-based binding proteins to different classes of targets.

The following examples are intended to illustrate but not limit the invention.

EXAMPLE I

Construction of the Fn3 gene

A synthetic gene for tenth Fn3 of fibronectin (Fig.1) was designed on the basis of amino acid residue 1416-1509 of human fibronectin (Kornblihtt, *et al.*, 1985) and its three dimensional structure (Main, *et al.*, 1992). The gene was engineered to include convenient restriction sites for mutagenesis and the so-called "preferred codons" for high level protein expression (Gribskov, *et al.*, 1984) were used. In addition, a glutamine residue was inserted after the N-terminal methionine in order to avoid partial processing of the N-terminal methionine which often degrades NMR spectra (Smith, *et al.*, 1994). Chemical reagents were of the analytical grade or better and purchased from Sigma Chemical Company and J.T. Baker, unless otherwise noted. Recombinant DNA procedures were performed as described in "Molecular Cloning" (Sambrook, *et al.*, 1989), unless otherwise stated. Custom oligonucleotides were purchased from Operon Technologies. Restriction and modification enzymes were from New England Biolabs.

The gene was assembled in the following manner. First, the gene sequence (Fig. 5) was divided into five parts with boundaries at designed restriction sites: fragment 1, NdeI-PstI (oligonucleotides FN1F and FN1R (Table 2); fragment 2, PstI-EcoRI (FN2F and FN2R); fragment 3, EcoRI-SalI (FN3F and FN3R); fragment 4, SalI-SacI (FN4F and FN4R); fragment 5, SacI-BamHI (FN5F and FN5R). Second, for each part, a pair of oligonucleotides which code opposite strands and have complementary overlaps of approximately 15 bases was synthesized. These oligonucleotides were designated FN1F-FN5R and are shown in Table 2. Third, each pair (e.g., FN1F and FN1R) was annealed and single-strand regions were filled in using the Klenow fragment of DNA polymerase. Fourth, the double stranded oligonucleotide was digested with the relevant restriction enzymes at the termini of the fragment and cloned into the pBlueScript SK plasmid (Stratagene) which had been digested with the same enzymes as those used for the fragments. The DNA sequence of the inserted fragment was confirmed by DNA sequencing using an

Applied Biosystems DNA sequencer and the dideoxy termination protocol provided by the manufacturer. Last, steps 2-4 were repeated to obtain the entire gene.

The gene was also cloned into the pET3a and pET15b (Novagen) vectors (pAS45 and pAS25, respectively). The maps of the plasmids are shown in Figs. 6 and 7. *E. coli* BL21 (DE3) (Novagen) containing these vectors expressed the Fn3 gene under the control of bacteriophage T7 promotor (Studier, *et al.*, 1990); pAS24 expresses the 96-residue Fn3 protein only, while pAS45 expresses Fn3 as a fusion protein with poly-histidine peptide (His•tag). High level expression of the Fn3 protein and its derivatives in *E. coli* was detected as an intense band on SDS-PAGE stained with CBB.

The binding reaction of the monobodies is characterized quantitatively by means of fluorescence spectroscopy using purified soluble monobodies.

Intrinsic fluorescence is monitored to measure binding reactions. Trp fluorescence (excitation at ~290 nm, emission at 300-350 nm) and Tyr fluorescence (excitation at ~260 nm, emission at ~303 nm) is monitored as the Fn3-mutant solution ($\leq 100 \mu\text{M}$) is titrated with a ligand solution. When a ligand is fluorescent (e.g. fluorescein), fluorescence from the ligand may be used. K_d of the reaction will be determined by the nonlinear least-squares fitting of the bimolecular binding equation.

If intrinsic fluorescence cannot be used to monitor the binding reaction, monobodies are labeled with fluorescein-NHS (Pierce) and fluorescence polarization is used to monitor the binding reaction (Burke *et al.*, 1996).

EXAMPLE II

Modifications to include restriction sites in the Fn3 gene

The restriction sites were incorporated in the synthetic Fn3 gene without changing the amino acid sequence Fn3. The positions of the restriction sites were chosen so that the gene construction could be completed without synthesizing long (>60 bases) oligonucleotides and so that two loop regions could be mutated

(including by randomization) by the cassette mutagenesis method (i.e., swapping a fragment with another synthetic fragment containing mutations). In addition, the restriction sites were chosen so that most sites were unique in the vector for phage display. Unique restriction sites allow one to recombine monobody clones which have been already selected in order to supply a larger sequence space.

EXAMPLE III

Construction of M13 phage display libraries

A vector for phage display, pAS38 (for its map, see Fig. 8) was constructed as follows. The XbaI-BamHI fragment of pET12a encoding the signal peptide of OmpT was cloned at the 5' end of the Fn3 gene. The C-terminal region (from the FN5F and FN5R oligonucleotides, see Table 2) of the Fn3 gene was replaced with a new fragment consisting of the FN5F and FN5R' oligonucleotides (Table 2) which introduced a MluI site and a linker sequence for making a fusion protein with the pIII protein of bacteriophage M13. A gene fragment coding the C-terminal domain of M13 pIII was prepared from the wild-type gene III of M13mp18 using PCR (Corey, *et al.*, 1993) and the fragment was inserted at the 3' end of the OmpT-Fn3 fusion gene using the MluI and HindIII sites.

Phages were produced and purified using a helper phage, M13K07, according to a standard method (Sambrook, *et al.*, 1989) except that phage particles were purified by a second polyethylene glycol precipitation. Successful display of Fn3 on fusion phages was confirmed by ELISA (Harlow & Lane, 1988) using an antibody against fibronectin (Sigma) and a custom anti-FN3 antibody (Cocalico Biologicals, PA, USA).

Example IV above. Other nucleic acid phage display libraries having variegation in the CD, DE or EF loop can be prepared by similar methods.

EXAMPLE VI

5 Loop variegations in the FG and BC loop

10 A nucleic acid phage display library having seven variegated residues (residues number 78-84) in the FG loop and five variegated residues (residue number 26-30) in the BC loop was prepared. Variegations in the BC loop were prepared by site-directed mutagenesis (Kunkel, *et al.*) using the BC3 oligonucleotide described in Table 1. Variegations in the FG loop were introduced using site-directed mutagenesis using the BC loop library as the starting material, thereby resulting in libraries containing variegations in both BC and FG loops. The oligonucleotide FG2 has variegating residues 78-84 and oligonucleotide FG4 has variegating residues 77-81 and a deletion of residues 82-84.

15 A nucleic acid phage display library having five variegated residues (residues 78-84) in the FG loop and a three residue deletion (residues 82-84) in the FG loop, and five variegated residues (residues 26-30) in the BC loop, was prepared. The shorter FG loop was made in an attempt to reduce the flexibility of the FG loop; the loop was shown to be highly flexible in Fn3 by the NMR studies of Main, *et al.* (1992). A highly flexible loop may be disadvantageous to forming a binding site with a high affinity (a large entropy loss is expected upon the ligand binding, because the flexible loop should become more rigid). In addition, other Fn3 domains (besides human) have shorter FG loops (for sequence alignment, see Figure 12 in Dickinson, *et al.* (1994)).

20 Randomization was achieved by the use of oligonucleotides containing degenerate nucleotide sequence (oligonucleotide BC3 for variegating the BC loop and oligonucleotides FG2 and FG4 for variegating the FG loops).

Site-directed mutagenesis was performed following published methods (see for example, Kunkel, 1985). The libraries were constructed by electrotransforming

09637614-03100
5 *E. coli* XL-1 Blue (Stratagene). Typically a library contains 10^8 to 10^9 independent clones. Library 2 contains five variegated residues in the BC loop and seven variegated residues in the FG loop. Library 4 contains five variegated residues in each of the BC and FG loops, and the length of the FG loop was shortened by three residues.

EXAMPLE VII

fd phage display libraries constructed with loop variegations

10 Phage display libraries are constructed using the fd phage as the genetic vector. The Fn3 gene is inserted in fUSE5 (Parmley & Smith, 1988) using SfiI restriction sites which are introduced at the 5' and 3' ends of the Fn3 gene using PCR. The expression of this phage results in the display of the fusion pIII protein on the surface of the fd phage. Variegations in the Fn3 loops are introduced using site-directed mutagenesis as described hereinabove, or by subcloning the Fn3
15 libraries constructed in M13 phage into the fUSE5 vector.

EXAMPLE VIII

Other phage display libraries

20 T7 phage libraries (Novagen, Madison, WI) and bacterial pili expression systems (Invitrogen) are also useful to express the Fn3 gene.

EXAMPLE IX

Isolation of polypeptides which bind to macromolecular structures

25 The selection of phage-displayed monobodies was performed following the protocols of Barbas and coworkers (Rosenblum & Barbas, 1995). Briefly, approximately 1 μ g of a target molecule ("antigen") in sodium carbonate buffer (100 mM, pH 8.5) was immobilized in the wells of a microtiter plate (Maxisorp, Nunc) by incubating overnight at 4°C in an air tight container. After the removal of this solution, the wells were then blocked with a 3% solution of BSA (Sigma, Fraction

OTEB"4T9E960

V) in TBS by incubating the plate at 37°C for 1 hour. A phagemid library solution (50 µl) containing approximately 10¹² colony forming units (cfu) of phagemid was absorbed in each well at 37°C for 1 hour. The wells were then washed with an appropriate buffer (typically TBST, 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, and 0.5% Tween20) three times (once for the first round). Bound phage were eluted by an acidic solution (typically, 0.1 M glycine-HCl, pH 2.2; 50 µl) and recovered phage were immediately neutralized with 3 µl of Tris solution. Alternatively, bound phage were eluted by incubating the wells with 50 µl of TBS containing the antigen (1 - 10 µM). Recovered phage were amplified using the standard protocol employing the XL1Blue cells as the host (Sambrook, *et al.*). The selection process was repeated 5-6 times to concentrate positive clones. After the final round, individual clones were picked and their binding affinities and DNA sequences were determined.

The binding affinities of monobodies on the phage surface were characterized using the phage ELISA technique (Li, *et al.*, 1995). Wells of microtiter plates (Nunc) were coated with an antigen and blocked with BSA. Purified phages (10⁸- 10¹¹ cfu) originating from a single colony were added to each well and incubated 2 hours at 37°C. After washing wells with an appropriate buffer (see above), bound phage were detected by the standard ELISA protocol using anti-M13 antibody (rabbit, Sigma) and anti-rabbit Ig-peroxidase conjugate (Pierce). Colorimetric assays were performed using Turbo-TMB (3,3',5,5'-tetramethylbenzidine, Pierce) as a substrate.

The binding affinities of monobodies on the phage surface were further characterized using the competition ELISA method (Djavadi-Ohanian, *et al.*, 1996). In this experiment, phage ELISA is performed in the same manner as described above, except that the phage solution contains a ligand at varied concentrations. The phage solution was incubated at 4°C for one hour prior to the binding of an immobilized ligand in a microtiter plate well. The affinities of phage

displayed monobodies are estimated by the decrease in ELISA signal as the free ligand concentration is increased.

After preliminary characterization of monobodies displayed on the surface of phage using phage ELISA, genes for positive clones were subcloned into the expression vector pAS45. *E. coli* BL21(DE3) (Novagen) was transformed with an expression vector (pAS45 and its derivatives). Cells were grown in M9 minimal medium and M9 medium supplemented with Bactotryptone (Difco) containing ampicillin (200 µg/ml). For isotopic labeling, ^{15}N NH_4Cl and/or ^{13}C glucose replaced unlabeled components. Stable isotopes were purchased from Isotec and Cambridge Isotope Labs. 500 ml medium in a 2 l baffie flask was inoculated with 10 ml of overnight culture and agitated at approximately 140 rpm at 37°C. IPTG was added at a final concentration of 1 mM to induce protein expression when OD(600 nm) reached approximately 1.0. The cells were harvested by centrifugation 3 hours after the addition of IPTG and kept frozen at -70°C until used.

Fn3 and monobodies with His•tag were purified as follows. Cells were suspended in 5 ml/(g cell) of 50 mM Tris (pH 7.6) containing 1 mM phenylmethylsulfonyl fluoride. HEL (Sigma, 3X crystallized) was added to a final concentration of 0.5 mg/ml. After incubating the solution for 30 min at 37°C, it was sonicated so as to cause cell breakage three times for 30 seconds on ice. Cell debris was removed by centrifugation at 15,000 rpm in an Sorval RC-2B centrifuge using an SS-34 rotor. Concentrated sodium chloride is added to the solution to a final concentration of 0.5 M. The solution was then applied to a 1 ml HisTrap™ chelating column (Pharmacia) preloaded with nickel chloride (0.1 M, 1 ml) and equilibrated in the Tris buffer (50 mM, pH 8.0) containing 0.5 M sodium chloride. After washing the column with the buffer, the bound protein was eluted with a Tris buffer (50 mM, pH 8.0) containing 0.5 M imidazole. The His•tag portion was cleaved off, when required, by treating the fusion protein with thrombin using the protocol supplied by Novagen (Madison, WI). Fn3 was separated from the His•tag

peptide and thrombin by a Resources[®] column (Pharmacia) using a linear gradient of sodium chloride (0 - 0.5 M) in sodium acetate buffer (20 mM, pH 5.0).

Small amounts of soluble monobodies were prepared as follows. XL-1 Blue cells containing pAS38 derivatives (plasmids coding Fn3-pIII fusion proteins) were grown in LB media at 37°C with vigorous shaking until OD(600 nm) reached approximately 1.0; IPTG was added to the culture to a final concentration of 1 mM, and the cells were further grown overnight at 37°C. Cells were removed from the medium by centrifugation, and the supernatant was applied to a microtiter well coated with a ligand. Although XL-1 Blue cells containing pAS38 and its derivatives express FN3-pIII fusion proteins, soluble proteins are also produced due to the cleavage of the linker between the Fn3 and pIII regions by proteolytic activities of *E. coli* (Rosenblum & Barbas, 1995). Binding of a monobody to the ligand was examined by the standard ELISA protocol using a custom antibody against Fn3 (purchased from Cocalico Biologicals, Reamstown, PA). Soluble monobodies obtained from the periplasmic fraction of *E. coli* cells using a standard osmotic shock method were also used.

EXAMPLE X

Ubiquitin binding monobody

Ubiquitin is a small (76 residue) protein involved in the degradation pathway in eukaryotes. It is a single domain globular protein. Yeast ubiquitin was purchased from Sigma Chemical Company and was used without further purification.

Libraries 2 and 4, described in Example VI above, were used to select ubiquitin-binding monobodies. Ubiquitin (1 µg in 50 µl sodium bicarbonate buffer (100 mM, pH 8.5)) was immobilized in the wells of a microtiter plate, followed by blocking with BSA (3% in TBS). Panning was performed as described above. In the first two rounds, 1 µg of ubiquitin was immobilized per well, and bound phage were eluted with an acidic solution. From the third to the sixth rounds, 0.1 µg of

ubiquitin was immobilized per well and the phage were eluted either with an acidic solution or with TBS containing 10 μ M ubiquitin.

Binding of selected clones was tested first in the polyclonal mode, i.e., before isolating individual clones. Selected clones from all libraries showed significant binding to ubiquitin. These results are shown in Figure 9. The binding to the immobilized ubiquitin of the clones was inhibited almost completely by less than 30 μ M soluble ubiquitin in the competition ELISA experiments (see Fig. 10). The sequences of the BC and FG loops of ubiquitin-binding monobodies is shown in Table 3.

10

Table 3. Sequences of ubiquitin-binding monobodies

	<u>Name</u>	<u>BC loop</u>	<u>FG loop</u>	<u>Occurrence (if more than one)</u>
15	211	CARRA	RWIPLAK	2
	212	CWRRRA	RWVGLAW	
	213	CKHRR	FADLWWR	
	214	CRRGR	RGFMWLS	
	215	CNWRR	RAYRYRW	
20	411	SRLRR	PPWRV	9
	422	ARWTL	RRWWW	
	424	GQRTF	RRWWA	

The 411 clone, which was the most enriched clone, was characterized using phage ELISA. The 411 clone showed selective binding and inhibition of binding in the presence of about 10 μ M ubiquitin in solution (Fig. 11).

25

EXAMPLE XI

Methods for the immobilization of small molecules

Target molecules were immobilized in wells of a microtiter plate (Maxisorp, Nunc) as described hereinbelow, and the wells were blocked with BSA. In addition

30

to the use of carrier protein as described below, a conjugate of a target molecule in biotin can be made. The biotinylated ligand can then be immobilized to a microtiter plate well which has been coated with streptavidin.

In addition to the use of a carrier protein as described below, one could make a conjugate of a target molecule and biotin (Pierce) and immobilize a biotinylated ligand to a microtiter plate well which has been coated with streptavidin (Smith and Scott, 1993).

Small molecules may be conjugated with a carrier protein such as bovine serum albumin (BSA, Sigma), and passively adsorbed to the microtiter plate well. Alternatively, methods of chemical conjugation can also be used. In addition, solid supports other than microtiter plates can readily be employed.

EXAMPLE XII

Fluorescein binding monobody

Fluorescein has been used as a target for the selection of antibodies from combinatorial libraries (Barbas, *et al.* 1992). NHS-fluorescein was obtained from Pierce and used according to the manufacturer's instructions in preparing conjugates with BSA (Sigma). Two types of fluorescein-BSA conjugates were prepared with approximate molar ratios of 17 (fluorescein) to one (BSA).

The selection process was repeated 5-6 times to concentrate positive clones. In this experiment, the phage library was incubated with a protein mixture (BSA, cytochrome C (Sigma, Horse) and RNaseA (Sigma, Bovine), 1 mg/ml each) at room temperature for 30 minutes, prior to the addition to ligand coated wells. Bound phage were eluted in TBS containing 10 μ M soluble fluorescein, instead of acid elution. After the final round, individual clones were picked and their binding affinities (see below) and DNA sequences were determined.

Table 4. Clones from Library #2

		<u>BC</u>	<u>FG</u>
	WT	AVTVR	RGDSPAS
5			
	pLB24.1	CNWRR	RAYRYRW
	pLB24.2	CMWRA	RWGMLRR
	pLB24.3	ARMRE	RWLRGRY
	pLB24.4	CARRR	RRAGWGW
10	pLB24.5	CNWRR	RAYRYRW
	pLB24.6	RWRER	RHPWTER
	pLB24.7	CNWRR	RAYRYRW
	pLB24.8	ERRVP	RLLWQR
	pLB24.9	GRGAG	FGSFERR
15	pLB24.11	CRWTR	RRWFDGA
	pLB24.12	CNWRR	RAYRYRW

Clones from Library #4

20	WT	AVTVR	GRGDS
	pLB25.1	GQRTF	RRWWA
	pLB25.2	GQRTF	RRWWA
	pLB25.3	GQRTF	RRWWA
25	pLB25.4	LRYRS	GWRWR
	pLB25.5	GQRTF	RRWWA
	pLB25.6	GQRTF	RRWWA
	pLB25.7	LRYRS	GWRWR
	pLB25.9	LRYRS	GWRWR
30	pLB25.11	GQRTF	RRWWA
	pLB25.12	LRYRS	GWRWR

Preliminary characterization of the binding affinities of selected clones were performed using phage ELISA and competition phage ELISA (see Fig. 12 (Fluorescein-1) and Fig. 13 (Fluorescein-2)). The four clones tested showed specific binding to the ligand-coated wells, and the binding reactions are inhibited by soluble
5 fluorescein (see Fig. 13).

EXAMPLE XIII

Digoxigenin binding monobody

Digoxigenin-3-O-methyl-carbonyl-e-aminocaproic acid-NHS (Boehringer
10 Mannheim) is used to prepare a digoxigenin-BSA conjugate. The coupling reaction is performed following the manufacturers' instructions. The digoxigenin-BSA conjugate is immobilized in the wells of a microtiter plate and used for panning. Panning is repeated 5 to 6 times to enrich binding clones. Because digoxigenin is sparingly soluble in aqueous solution, bound phages are eluted from the well using
15 acidic solution. See Example XIV.

EXAMPLE XIV

TSAC (transition state analog compound) binding monobodies

Carbonate hydrolyzing monobodies are selected as follows. A transition
20 state analog for carbonate hydrolysis, 4-nitrophenyl phosphonate is synthesized by an Arbuzov reaction as described previously (Jacobs and Schultz, 1987). The phosphonate is then coupled to the carrier protein, BSA, using carbodiimide, followed by exhaustive dialysis (Jacobs and Schultz, 1987). The hapten-BSA conjugate is immobilized in the wells of a microtiter plate and monobody selection
25 is performed as described above. Catalytic activities of selected monobodies are tested using 4-nitrophenyl carbonate as the substrate.

Other haptens useful to produce catalytic monobodies are summarized in H. Suzuki (1994) and in N. R. Thomas (1994).

EXAMPLE XV

NMR characterization of Fn3 and comparison of the Fn3 secreted by yeast with that secreted by *E. coli*

Nuclear magnetic resonance (NMR) experiments are performed to identify
5 the contact surface between FnAb and a target molecule, e.g., monoclonal antibodies to
fluorescein, ubiquitin, RNaseA and soluble derivatives of digoxigenin. The
information is then be used to improve the affinity and specificity of the monoclonal antibody.
Purified monoclonal antibody samples are dissolved in an appropriate buffer for NMR
spectroscopy using Amicon ultrafiltration cell with a YM-3 membrane. Buffers are
10 made with 90 % H₂O/10 % D₂O (distilled grade, Isotec) or with 100 % D₂O.
Deuterated compounds (e.g. acetate) are used to eliminate strong signals from them.

NMR experiments are performed on a Varian Unity INOVA 600
spectrometer equipped with four RF channels and a triple resonance probe with
pulsed field gradient capability. NMR spectra are analyzed using processing
15 programs such as Felix (Molecular Simulations), nmrPipe, PIPP, and CAPP
(Garrett, *et al.*, 1991; Delaglio, *et al.*, 1995) on UNIX workstations. Sequence
specific resonance assignments are made using well-established strategy using a set
of triple resonance experiments (CBCA(CO)NH and HNCACB) (Grzesiek & Bax,
1992; Wittenkind & Mueller, 1993).

20 Nuclear Overhauser effect (NOE) is observed between ¹H nuclei closer than
approximately 5 Å, which allows one to obtain information on interproton distances.
A series of double- and triple-resonance experiments (Table 5; for recent reviews on
these techniques, see Bax & Grzesiek, 1993 and Kay, 1995) are performed to collect
distance (i.e. NOE) and dihedral angle (J-coupling) constraints. Isotope-filtered
25 experiments are performed to determine resonance assignments of the bound ligand
and to obtain distance constraints within the ligand and those between FnAb and the
ligand. Details of sequence specific resonance assignments and NOE peak
assignments have been described in detail elsewhere (Clore & Gronenborn, 1991;
Pascal, *et al.*, 1994b; Metzler, *et al.*, 1996).

Table 5. NMR experiments for structure characterization

	<u>Experiment Name</u>	<u>Reference</u>
	1. reference spectra	
5	2D- ¹ H, ¹⁵ N-HSQC	(Bodenhausen & Ruben, 1980; Kay, <i>et al.</i> , 1992)
	2D- ¹ H, ¹³ C-HSQC	(Bodenhausen & Ruben, 1980; Vuister & Bax, 1992)
	2. backbone and side chain resonance assignments of ¹³ C/ ¹⁵ N-labeled protein	
10	3D-CBCA(CO)NH	(Grzesiek & Bax, 1992)
	3D-HNCACB	(Wittenkind & Mueller, 1993)
	3D-C(CO)NH	(Logan <i>et al.</i> , 1992; Grzesiek <i>et al.</i> , 1993)
	3D-H(CCO)NH	
15	3D-HBHA(CBCACO)NH	(Grzesiek & Bax, 1993)
	3D-HCCH-TOCSY	(Kay <i>et al.</i> , 1993)
	3D-HCCH-COSY	(Ikura <i>et al.</i> , 1991)
	3D- ¹ H, ¹⁵ N-TOCSY-HSQC	(Zhang <i>et al.</i> , 1994)
	2D-HB(CBCDCE)HE	(Yamazaki <i>et al.</i> , 1993)
20	3. resonance assignments of unlabeled ligand	
	2D-isotope-filtered ¹ H-TOCSY	
	2D-isotope-filtered ¹ H-COSY	
25	2D-isotope-filtered ¹ H-NOESY	(Ikura & Bax, 1992)
	4. structural constraints	
	<i>within labeled protein</i>	
	3D- ¹ H, ¹⁵ N-NOESY-HSQC	(Zhang <i>et al.</i> , 1994)
30	4D- ¹ H, ¹³ C-HMQC-NOESY-HMQC	(Vuister <i>et al.</i> , 1993)
	4D- ¹ H, ¹³ C, ¹⁵ N-HSQC-NOESY-HSQC	(Muhandiram <i>et al.</i> , 1993; Pascal <i>et al.</i> , 1994a)

within unlabeled ligand

2D-isotope-filtered ^1H -NOESY (Ikura & Bax, 1992)

interactions between protein and ligand

3D-isotope-filtered ^1H , ^{15}N -NOESY-HSQC

5 3D-isotope-filtered ^1H , ^{13}C -NOESY-HSQC (Lee *et al.*, 1994)

5. dihedral angle constraints

J-modulated ^1H , ^{15}N -HSQC (Billeter *et al.*, 1992)

10 3D-HNHB (Archer *et al.*, 1991)

Backbone ^1H , ^{15}N and ^{13}C resonance assignments for a monobody are compared to those for wild-type Fn3 to assess structural changes in the mutant.

15 Once these data establish that the mutant retains the global structure, structural refinement is performed using experimental NOE data. Because the structural difference of a monobody is expected to be minor, the wild-type structure can be used as the initial model after modifying the amino acid sequence. The mutations are introduced to the wild-type structure by interactive molecular modeling, and then
20 the structure is energy-minimized using a molecular modeling program such as Quanta (Molecular Simulations). Solution structure is refined using cycles of dynamical simulated annealing (Nilges *et al.*, 1988) in the program X-PLOR (Brünger, 1992). Typically, an ensemble of fifty structures is calculated. The validity of the refined structures is confirmed by calculating a fewer number of
25 structures from randomly generated initial structures in X-PLOR using the YASAP protocol (Nilges, *et al.*, 1991). Structure of a monobody-ligand complex is calculated by first refining both components individually using intramolecular NOEs, and then docking the two using intermolecular NOEs.

For example, the ^1H , ^{15}N -HSQC spectrum for the fluorescein-binding
30 monobody LB25.5 is shown in Figure 14. The spectrum shows a good dispersion

(peaks are spread out) indicating that LB25.5 is folded into a globular conformation. Further, the spectrum resembles that for the wild-type Fn3, showing that the overall structure of LB25.5 is similar to that of Fn3. These results demonstrate that ligand-binding monobodies can be obtained without changing the global fold of the Fn3 scaffold.

Chemical shift perturbation experiments are performed by forming the complex between an isotope-labeled FnAb and an unlabeled ligand. The formation of a stoichiometric complex is followed by recording the HSQC spectrum. Because chemical shift is extremely sensitive to nuclear environment, formation of a complex usually results in substantial chemical shift changes for resonances of amino acid residues in the interface. Isotope-edited NMR experiments (2D HSQC and 3D CBCA(CO)NH) are used to identify the resonances that are perturbed in the labeled component of the complex; i.e. the monobody. Although the possibility of artifacts due to long-range conformational changes must always be considered, substantial differences for residues clustered on continuous surfaces are most likely to arise from direct contacts (Chen *et al.*, 1993; Gronenborn & Clore, 1993).

An alternative method for mapping the interaction surface utilizes amide hydrogen exchange (HX) measurements. HX rates for each amide proton are measured for ^{15}N labeled monobody both free and complexed with a ligand. Ligand binding is expected to result in decreased amide HX rates for monobody residues in the interface between the two proteins, thus identifying the binding surface. HX rates for monobodies in the complex are measured by allowing HX to occur for a variable time following transfer of the complex to D_2O ; the complex is dissociated by lowering pH and the HSQC spectrum is recorded at low pH where amide HX is slow. Fn3 is stable and soluble at low pH, satisfying the prerequisite for the experiments.

EXAMPLE XVI

Construction and Analysis of Fn3-Display System Specific for Ubiquitin

An Fn3-display system was designed and synthesized, ubiquitin-binding clones were isolated and a major Fn3 mutant in these clones was biophysically characterized.

Gene construction and phage display of Fn3 was performed as in Examples I and II above. The Fn3-phage pIII fusion protein was expressed from a phagemid-display vector, while the other components of the M13 phage, including the wild-type pIII, were produced using a helper phage (Bass et al., 1990). Thus, a phage produced by this system should contain less than one copy of Fn3 displayed on the surface. The surface display of Fn3 on the phage was detected by ELISA using an anti-Fn3 antibody. Only phages containing the Fn3-pIII fusion vector reacted with the antibody.

After confirming the phage surface to display Fn3, a phage display library of Fn3 was constructed as in Example III. Random sequences were introduced in the BC and FG loops. In the first library, five residues (77-81) were randomized and three residues (82-84) were deleted from the FG loop. The deletion was intended to reduce the flexibility and improve the binding affinity of the FG loop. Five residues (26-30) were also randomized in the BC loop in order to provide a larger contact surface with the target molecule. Thus, the resulting library contains five randomized residues in each of the BC and FG loops (Table 6). This library contained approximately 10^8 independent clones.

Library Screening

Library screening was performed using ubiquitin as the target molecule. In each round of panning, Fn3-phages were absorbed to a ubiquitin-coated surface, and bound phages were eluted competitively with soluble ubiquitin. The recovery ratio improved from 4.3×10^{-7} in the second round to 4.5×10^{-6} in the fifth round, suggesting an enrichment of binding clones. After five rounds of panning, the amino acid sequences of individual clones were determined (Table 6).

Table 6. Sequences in the variegated loops of enriched clones

Name	BC loop	FG loop	Frequency
Wild Type	GCAGTTACCGTGCGT	GGCCGTGGTGACAGCCCAGCGAGC	–
	AlaValThrValArg	GlyArgGlyAspSerProAlaSer	
Library ^a	NNKNNKNNKNNKNNK	NNKNNKNNKNNKNNK-----	–
	X X X X X	X X X X X (deletion)	
5 clone1	TCGAGGTTGCGGCGG	CCGCCGTGGAGGGTG	9
(Ubi4)	SerArgLeuArgArg	ProProTrpArgVal	
clone2	GGTCAGCGAACTTTT	AGGCGGTGGTGGGCT	1
	GlyGlnArgThrPhe	ArgArgTrpTrpAla	
clone3	GCGAGGTGGACGCTT	AGGCGGTGGTGGTGG	1
	AlaArgTrpThrLeu	ArgArgTrpTrpTrp	

10 ^a N denotes an equimolar mixture of A, T, G and C; K denotes an equimolar mixture of G and T.

A clone, dubbed Ubi4, dominated the enriched pool of Fn3 variants. Therefore, further investigation was focused on this Ubi4 clone. Ubi4 contains four mutations in the BC loop (Arg 30 in the BC loop was conserved) and five mutations and three
 15 deletions in the FG loop. Thus 13% (12 out of 94) of the residues were altered in Ubi4 from the wild-type sequence.

Figure 15 shows a phage ELISA analysis of Ubi4. The Ubi4 phage binds to the target molecule, ubiquitin, with a significant affinity, while a phage displaying the wild-type Fn3 domain or a phase with no displayed molecules show little
 20 detectable binding to ubiquitin (Figure 15a). In addition, the Ubi4 phage showed a somewhat elevated level of background binding to the control surface lacking the ubiquitin coating. A competition ELISA experiments shows the IC₅₀ (concentration of the free ligand which causes 50% inhibition of binding) of the binding reaction is approximately 5 μM (Fig. 15b). BSA, bovine ribonuclease A and cytochrome C
 25 show little inhibition of the Ubi4-ubiquitin binding reaction (Figure 15c), indicating that the binding reaction of Ubi4 to ubiquitin does result from specific binding.

Characterization of a Mutant Fn3 Protein

The expression system yielded 50-100 mg Fn3 protein per liter culture. A similar level of protein expression was observed for the Ubi4 clone and other mutant Fn3 proteins.

- 5 Ubi4-Fn3 was expressed as an independent protein. Though a majority of Ubi4 was expressed in *E. coli* as a soluble protein, its solubility was found to be significantly reduced as compared to that of wild-type Fn3. Ubi4 was soluble up to ~20 μ M at low pH, with much lower solubility at neutral pH. This solubility was not high enough for detailed structural characterization using NMR spectroscopy or
- 10 X-ray crystallography.

- The solubility of the Ubi4 protein was improved by adding a solubility tail, GKKKGK, as a C-terminal extension. The gene for Ubi4-Fn3 was subcloned into the expression vector pAS45 using PCR. The C-terminal solubilization tag, GKKKGK, was incorporated in this step. *E. coli* BL21 (DE3) (Novagen) was transformed with
- 15 the expression vector (pAS45 and its derivatives). Cells were grown in M9 minimal media and M9 media supplemented with Bactotryptone (Difco) containing ampicillin (200 μ g/ml). For isotopic labeling, 15 N NH_4Cl replaced unlabeled NH_4Cl in the media. 500 ml medium in a 2 liter baffle flask was inoculated with 10 ml of overnight culture and agitated at 37°C. IPTG was added at a final concentration of 1
- 20 mM to initiate protein expression when OD (600 nm) reaches one. The cells were harvested by centrifugation 3 hours after the addition of IPTG and kept frozen at -70°C until used.

- Proteins were purified as follows. Cells were suspended in 5 ml/(g cell) of Tris (50 mM, pH 7.6) containing phenylmethylsulfonyl fluoride (1 mM). Hen egg
- 25 lysozyme (Sigma) was added to a final concentration of 0.5 mg/ml. After incubating the solution for 30 minutes at 37°C, it was sonicated three times for 30 seconds on ice. Cell debris was removed by centrifugation. Concentrated sodium chloride was added to the solution to a final concentration of 0.5 M. The solution was applied to a Hi-Trap chelating column (Pharmacia) preloaded with nickel and equilibrated in

the Tris buffer containing sodium chloride (0.5 M). After washing the column with the buffer, histag-Fn3 was eluted with the buffer containing 500 mM imidazole. The protein was further purified using a ResourceS column (Pharmacia) with a NaCl gradient in a sodium acetate buffer (20 mM, pH 4.6).

- 5 With the GKKGK tail, the solubility of the Ubi4 protein was increased to over 1 mM at low pH and up to ~50 μ M at neutral pH. Therefore, further analyses were performed on Ubi4 with this C-terminal extension (hereafter referred to as Ubi4-K). It has been reported that the solubility of a minibody could be significantly improved by addition of three Lys residues at the N- or C-termini
- 10 (Bianchi et al., 1994). In the case of protein Rop, a non-structured C-terminal tail is critical in maintaining its solubility (Smith et al., 1995).

- Oligomerization states of the Ubi4 protein were determined using a size exclusion column. The wild-type Fn3 protein was monomeric at low and neutral pH's. However, the peak of the Ubi4-K protein was significantly broader than that
- 15 of wild-type Fn3, and eluted after the wild-type protein. This suggests interactions between Ubi4-K and the column material, precluding the use of size exclusion chromatography to determine the oligomerization state of Ubi4. NMR studies suggest that the protein is monomeric at low pH.

- The Ubi4-K protein retained a binding affinity to ubiquitin as judged by
- 20 ELISA (Figure 15d). However, an attempt to determine the dissociation constant using a biosensor (Affinity Sensors, Cambridge, U.K.) failed because of high background binding of Ubi4-K-Fn3 to the sensor matrix. This matrix mainly consists of dextran, consistent with our observation that interactions between Ubi4-K interacts with the cross-linked dextran of the size exclusion column.

25

Example XVII

Stability Measurements of Monobodies

Guanidine hydrochloride (GuHCl)-induced unfolding and refolding reactions were followed by measuring tryptophan fluorescence. Experiments were performed

on a Spectronic AB-2 spectrofluorometer equipped with a motor-driven syringe (Hamilton Co.). The cuvette temperature was kept at 30°C. The spectrofluorometer and the syringe were controlled by a single computer using a home-built interface. This system automatically records a series of spectra following GuHCl titration. An experiment started with a 1.5 ml buffer solution containing 5 μ M protein. An emission spectrum (300-400 nm; excitation at 290 nm) was recorded following a delay (3-5 minutes) after each injection (50 or 100 μ l) of a buffer solution containing GuHCl. These steps were repeated until the solution volume reached the full capacity of a cuvette (3.0 ml). Fluorescence intensities were normalized as ratios to the intensity at an isofluorescent point which was determined in separate experiments. Unfolding curves were fitted with a two-state model using a nonlinear least-squares routine (Santoro & Bolen, 1988). No significant differences were observed between experiments with delay times (between an injection and the start of spectrum acquisition) of 2 minutes and 10 minutes, indicating that the unfolding/refolding reactions reached close to an equilibrium at each concentration point within the delay times used.

Conformational stability of Ubi4-K was measured using above-described GuHCl-induced unfolding method. The measurements were performed under two sets of conditions; first at pH 3.3 in the presence of 300 mM sodium chloride, where Ubi4-K is highly soluble, and second in TBS, which was used for library screening. Under both conditions, the unfolding reaction was reversible, and we detected no signs of aggregation or irreversible unfolding. Figure 16 shows unfolding transitions of Ubi4-K and wild-type Fn3 with the N-terminal (his)₆ tag and the C-terminal solubility tag. The stability of wild-type Fn3 was not significantly affected by the addition of these tags. Parameters characterizing the unfolding transitions are listed in Table 7.

Table 7. Stability parameters for Ubi4 and wild-type Fn3 as determined by GuHCl-induced unfolding

	Protein	ΔG_0 (kcal mol ⁻¹)	m_G (kcal mol ⁻¹ M ⁻¹)
5	Ubi4 (pH 7.5)	4.8 ± 0.1	2.12 ± 0.04
	Ubi4 (pH 3.3)	6.5 ± 0.1	2.07 ± 0.02
	Wild-type (pH 7.5)	7.2 ± 0.2	1.60 ± 0.04
	Wild-type (pH 3.3)	11.2 ± 0.1	2.03 ± 0.02

- 10 ΔG_0 is the free energy of unfolding in the absence of denaturant; m_G is the dependence of the free energy of unfolding on GuHCl concentration. For solution conditions, see Figure 4 caption.

Though the introduced mutations in the two loops certainly decreased the stability of
 15 Ubi4-K relative to wild-type Fn3, the stability of Ubi4 remains comparable to that of a “typical” globular protein. It should also be noted that the stabilities of the wild-type and Ubi4-K proteins were higher at pH 3.3 than at pH 7.5.

The Ubi4 protein had a significantly reduced solubility as compared to that of wild-type Fn3, but the solubility was improved by the addition of a solubility tail.
 20 Since the two mutated loops comprise the only differences between the wild-type and Ubi4 proteins, these loops must be the origin of the reduced solubility. At this point, it is not clear whether the aggregation of Ubi4-K is caused by interactions between the loops, or by interactions between the loops and the invariable regions of the Fn3 scaffold.

25 The Ubi4-K protein retained the global fold of Fn3, showing that this scaffold can accommodate a large number of mutations in the two loops tested. Though the stability of the Ubi4-K protein is significantly lower than that of the wild-type Fn3 protein, the Ubi4 protein still has a conformational stability comparable to those for small globular proteins. The use of a highly stable domain
 30 as a scaffold is clearly advantageous for introducing mutations without affecting the global fold of the scaffold. In addition, the GuHCl-induced unfolding of the Ubi4 protein is almost completely reversible. This allows the preparation of a correctly folded protein even when a Fn3 mutant is expressed in a misfolded form, as in

inclusion bodies. The modest stability of Ubi4 in the conditions used for library screening indicates that Fn3 variants are folded on the phage surface. This suggests that a Fn3 clone is selected by its binding affinity in the folded form, not in a denatured form. Dickinson et al. proposed that Val 29 and Arg 30 in the BC loop stabilize Fn3. Val 29 makes contact with the hydrophobic core, and Arg 30 forms hydrogen bonds with Gly 52 and Val 75. In Ubi4-Fn3, Val 29 is replaced with Arg, while Arg 30 is conserved. The FG loop was also mutated in the library. This loop is flexible in the wild-type structure, and shows a large variation in length among human Fn3 domains (Main et al., 1992). These observations suggest that mutations in the FG loop may have less impact on stability. In addition, the N-terminal tail of Fn3 is adjacent to the molecular surface formed by the BC and FG loops (Figure 1 and 17) and does not form a well-defined structure. Mutations in the N-terminal tail would not be expected to have strong detrimental effects on stability. Thus, residues in the N-terminal tail may be good sites for introducing additional mutations.

Example XVIII

NMR Spectroscopy of Ubi4-Fn3

Ubi4-Fn3 was dissolved in [^2H]-Gly HCl buffer (20 mM, pH 3.3) containing NaCl (300 mM) using an Amicon ultrafiltration unit. The final protein concentration was 1 mM. NMR experiments were performed on a Varian Unity INOVA 600 spectrometer equipped with a triple-resonance probe with pulsed field gradient. The probe temperature was set at 30°C. HSQC, TOCSY-HSQC and NOESY-HSQC spectra were recorded using published procedures (Kay et al., 1992; Zhang et al., 1994). NMR spectra were processed and analyzed using the NMRPipe and NMRView software (Johnson & Blevins, 1994; Delaglio et al., 1995) on UNIX workstations. Sequence-specific resonance assignments were made using standard procedures (Wüthrich, 1986; Clore & Gronenborn, 1991). The assignments for wild-type Fn3 (Baron et al., 1992) were confirmed using a ^{15}N -labeled protein dissolved in sodium acetate buffer (50 mM, pH 4.6) at 30°C.

007130-192960

The three-dimensional structure of Ubi4-K was characterized using this heteronuclear NMR spectroscopy method. A high quality spectrum could be collected on a 1 mM solution of ^{15}N -labeled Ubi4 (Figure 17a) at low pH. The linewidth of amide peaks of Ubi4-K was similar to that of wild-type Fn3, suggesting that Ubi4-K is monomeric under the conditions used. Complete assignments for backbone ^1H and ^{15}N nuclei were achieved using standard ^1H , ^{15}N double resonance techniques, except for a row of His residues in the N-terminal (His)₆ tag. There were a few weak peaks in the HSQC spectrum which appeared to originate from a minor species containing the N-terminal Met residue. Mass spectroscopy analysis showed that a majority of Ubi4-K does not contain the N-terminal Met residue. Fig. 17 shows differences in ^1HN and ^{15}N chemical shifts between Ubi4-K and wild-type Fn3. Only small differences are observed in the chemical shifts, except for those in and near the mutated BC and FG loops. These results clearly indicate that Ubi4-K retains the global fold of Fn3, despite the extensive mutations in the two loops. A few residues in the N-terminal region, which is close to the two mutated loops, also exhibit significant chemical differences between the two proteins. An HSQC spectrum was also recorded on a 50 μM sample of Ubi4-K in TBS. The spectrum was similar to that collected at low pH, indicating that the global conformation of Ubi4 is maintained between pH 7.5 and 3.3.

20

The complete disclosure of all patents, patent documents and publications cited herein are incorporated by reference as if individually incorporated. The foregoing detailed description and examples have been given for clarity of understanding only. No unnecessary limitations are to be understood therefrom.

25 The invention is not limited to the exact details shown and described for variations obvious to one skilled in the art will be included within the invention defined by the claims.

REFERENCES

- Alzari, P.N., Lascombe, M.-B. & Poljak, R.J. (1988) Three-dimensional structure of antibodies. *Annu. Rev. Immunol.* 6, 555-580.
- Archer, S. J., Ikura, M., Torchia, D. A. & Bax, A. (1991) An alternative 3D NMR technique for correlating backbone ¹⁵N with side chain H β resonances in large proteins *J. Magn. Reson.* 95, 636-641.
- Aukhil, I., Joshi, P., Yan, Y. & Erickson, H. P. (1993) Cell- and heparin-binding domains of the hexabrachion arm identified by tenascin expression protein *J. Biol. Chem.* 268, 2542-2553.
- Barbas, C. F., III, Kang, A. S., Lerner, R. A. & Benkovic, S. J. (1991) Assembly of combinatorial antibody libraries on phage surfaces: the gene III site. *Proc. Natl. Acad. Sci. USA* 88, 7978-7982.
- Barbas, C. F., III, Bain, J. D., Hoekstra, D. M. & Lerner, R. A. (1992) Semisynthetic combinatorial libraries: A chemical solution to the diversity problem *Proc. Natl. Acad. Sci. USA* 89, 4457-4461.
- Baron, M., Main, A. L., Driscoll, P. C., Mardon, H. J., Boyd, J. & Campbell, I.D. (1992) ¹H NMR assignment and secondary structure of the cell adhesion type II module of fibronectin *Biochemistry* 31, 2068-2073.
- Baron, M., Norman, D. G. & Campbell, I. D. (1991) Protein modules *Trends Biochem. Sci.* 16, 13-17.
- Bass, S., Greene, R. & Wells, J. A. (1990) Hormone phage: An enrichment method for variant proteins with altered binding properties *Proteins: Struct. Funct. Genet.* 8, 309-314.
- Bax, A. & Grzesiek, S. (1993) Methodological advances in protein NMR. *Acc. Chem. Res.* 26, 131-138.
- Becktel, W. J. & Schellman, J. A. (1987) Protein stability curves. *Biopolymer* 26, 1859-1877.
- Bhat, T. N., Bentley, G. A., Boulot, G., Greene, M. I., Tello, D., Dall'acqua, W., Souchon, H., Schwarz, F. P., Mariuzza, R. A. & Poljak, R. J. (1994) Bound

- water molecules and conformational stabilization help mediate an antigen-antibody association. *Proc. Natl. Acad. Sci. USA* 91, 1089-1093.
- Bianchi, E., Venturini, S., Pessi, A., Tramontano, A. & Sollazzo, M. (1994) High level expression and rational mutagenesis of a designed protein, the minibody. From an insoluble to a soluble molecule. *J. Mol. Biol.* 236, 649-659.
- Billeter, M., Neri, D., Otting, G., Qian, Y. Q. & Wüthrich, K. (1992) Precise vicinal coupling constants $^3J_{\text{HNa}}$ in proteins from nonlinear fits of J-modulated [^{15}N , ^1H]-COSY experiments. *J. Biomol. NMR* 2, 257-274.
- Bodenhausen, G. & Ruben, D. J. (1980) Natural abundance nitrogen-15 NMR by enhanced heteronuclear spectroscopy. *Chem. Phys. Lett.* 69, 185-189.
- Bork, P. & Doolittle, R. F., *PNAS* 89:8990-8994 (1992).
- Bork, P. & Doolittle, R. F. (1992) Proposed acquisition of an animal protein domain by bacteria. *Proc. Natl. Acad. Sci. USA* 89, 8990-8994.
- Bork, P., Hom, L. & Sander, C. (1994) The immunoglobulin fold. Structural classification, sequence patterns and common core. *J. Mol. Biol.* 242, 309-320.
- Brünger, A. T. (1992) *X-PLOR (Version 3.1): A system for X-ray crystallography and NMR.*, Yale Univ. Press, New Haven.
- Burke, T., Bolger, R., Checovich, W. & Lowery, R. (1996) in *Phage display of peptides and proteins* (Kay, B. K., Winter, J. and McCafferty, J., Ed.) Vol. pp305-326, Academic Press, San Diego.
- Campbell, I. D. & Spitzfaden, C. (1994) Building proteins with fibronectin type III modules *Structure* 2, 233-337.
- Chen, Y., Reizer, J., Saier, M. H., Fairbrother, W. J. & Wright, P. E. (1993) Mapping the binding interfaces of the proteins of the bacterial phosphotransferase system, HPr and ΠAglc . *Biochemistry* 32, 32-37.
- Clackson & Wells, (1994) *Trends Biotechnology* 12, 173-184.

- Clore, G. M. & Gronenborn, A. M. (1991) Structure of larger proteins in solution: Three- and four-dimensional heteronuclear NMR spectroscopy. *Science* 252, 1390-1399.
- Corey, D. R., Shiau, A. K., Q., Y., Janowski, B. A. & Craik, C. S. (1993) Trypsin display on the surface of bacteriophage. *Gene* 128, 129-134.
- Davies, J. & Riechmann, L. (1996). Single antibody domains as small recognition units: design and *in vitro* antigen selection of camelized, human VH domains with improved protein stability. *Protein Eng.*, 9(6), 531-537.
- Davies, J. & Riechmann, L. (1995) Antibody VH domains as small recognition units. *Bio/Technol.* 13, 475-479.
- Delaglio, F., Grzesiek, S., Vuister, G. W., Zhu, G., Pfeifer, J. & Bax, A. (1995) NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J. Biomol. NMR* 6, 277-293.
- Deng, W. P. & Nickoloff, J. A. (1992) Site-directed mutagenesis of virtually any plasmid by eliminating a unique site. *Anal. Biochem.* 200, 81-88.
- deVos, A. M., Ultsch, M. & Kossiakoff, A. A. (1992) Human Growth hormone and extracellular domain of its receptor: crystal structure of the complex. *Science* 255, 306-312.
- Dickinson, C. D., Veerapandian, B., Dai, X.P., Hamlin, R. C., Xuong, N.-H., Ruoslahti, E. & Ely, K. R. (1994) Crystal structure of the tenth type III cell adhesion module of human fibronectin *J. Mol. Biol.* 236, 1079-1092.
- Djavadi-Ohanian, L., Goldberg, M. E. & Friguet, B. (1996) in *Antibody Engineering. A Practical Approach* (McCafferty, J., Hoogenboom, H. R. and Chiswell, D. J., Ed.) Vol. pp77-97, Oxford Univ. Press, Oxford.
- Dougall, W. C., Peterson, N. C. & Greene, M. I. (1994) Antibody-structure-based design of pharmacological agents. *Trends Biotechnol.* 12, 372-379.
- Garrett, D. S., Powers, R., Gronenborn, A. M. & Clore, G. M. (1991) A common sense approach to peak picking in two-, three- and four-dimensional spectra

using automatic computer analysis of contour diagrams. *J. Magn. Reson.* 95, 214-220.

Ghosh, G., Van Duyne, G., Ghosh, S. & Sigler, P. B. (1995) Structure of NF-kB p50 homodimer bound to a kB site. *Nature* 373, 303-310.

Gribskov, M., Devereux, J. & Burgess, R. R. (1984) The codon preference plot: graphic analysis of protein coding sequences and prediction of gene expression. *Nuc. Acids. Res.* 12, 539-549.

Groneborn, A. M., Filpula, D. R., Essig, N. Z., Achari, A., Whitlow, M., Wingfield, P. T. & Clore, G. M. (1991) A novel, highly stable fold of the immunoglobulin binding domain of Streptococcal protein G. *Science* 253, 657-661.

Gronenborn, A. M. & Clore, G. M. (1993) Identification of the contact surface of a Streptococcal protein G domain complexed with a human Fc fragment. *J. Mol. Biol.* 233, 331-335.

Grzesiek, S., Anglister, J. & Bax, A. (1993) Correlation of backbone amide and aliphatic side-chain resonances in ¹³C/¹⁵N-enriched proteins by isotropic mixing of ¹³C magnetization. *J. Magn. Reson. B* 101, 114-119.

Grzesiek, S. & Bax, A. (1992) Correlating backbone amide and side chain resonances in larger proteins by multiple relayed triple resonance NMR. *J. Am. Chem. Soc.* 114, 6291-6293.

Grzesiek, S. & Bax, A. (1993) Amino acid type determination in the sequential assignment procedure of uniformly ¹³C/¹⁵N-enriched proteins. *J. Biomol. NMR* 3, 185-204.

Harlow, E. & Lane, D. (1988) *Antibodies. A laboratory manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor.

Harpez, Y. & Chothia, C. (1994) Many of the immunoglobulin superfamily domains in cell adhesion molecules and surface receptors belong to a new structural set which is close to that containing variable domains *J. Mol. Biol.* 238, 528-539.

- Hawkins, R. E., Russell, S. J., Baier, M. & Winter, G. (1993) The contribution of contact and non-contact residues of antibody in the affinity of binding to antigen. The interaction of mutant D1.3 antibodies with lysozyme. *J. Mol. Biol.* 234, 958-964.
- Holliger, P. et al., (1993) *Proc. Natl. Acad. Sci.* 90, 6444-6448.
- Hu, S.-z., et al., *Cancer Res.* 56:3055-3061 (1996).
- Ikura, M. & Bax, A. (1992) Isotope-filtered 2D NMR of a protein-peptide complex: study of a skeletal muscle myosin light chain kinase fragment bound to calmodulin. *J. Am. Chem. Soc.* 114, 2433-2440.
- Ikura, M., Kay, L. E. & Bax, A. (1991) Improved three-dimensional ^1H - ^{13}C - ^1H correlation spectroscopy of a ^{13}C -labeled protein using constant-time evolution. *J. Biomol. NMR* 1, 299-304.
- Jacobs, J. & Schultz, P. G. (1987) Catalytic antibodies. *J. Am. Chem. Soc.* 109, 2174-2176.
- Janda, K. D., et al., *Science* 275:945-948 (1997).
- Jones, E. Y. (1993) The immunoglobulin superfamily *Curr. Opinion struct. Biol.* 3, 846-852.
- Jones, P. T., Dear, P. H., Foote, J., Neuberger, M. S. & Winter, G. (1986) Replacing the complementarity-determining regions in a human antibody with those from a mouse *Nature* 321, 522-525.
- Kabsch, W. & Sander, C. (1983) Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* 22, 2577-2637.
- Kay, L. E. (1995) Field gradient techniques in NMR spectroscopy. *Curr. Opinion Struct. Biol.* 5, 674-681.
- Kay, L. E., Keifer, P. & Saarinen, T. (1992) Pure absorption gradient enhanced heteronuclear single quantum correlation spectroscopy with improved sensitivity *J. Am. Chem. Soc.* 114, 10663-10665.

- Kay, L. E., Xu, G.-Y. & Singer, A. U. (1993) A Gradient-Enhanced HCCH-TOCSY Experiment for Recording Side-Chain ¹H and ¹³C Correlations in H₂O Samples of Proteins. *J. Magn. Reson. B* 101, 333-337.
- Koide, S., Dyson, H. J. & Wright, P. E. (1993) Characterization of a folding intermediate of apoplastcyanin trapped by proline isomerization. *Biochemistry* 32, 12299-12310.
- Kornblihtt, A. R., Umezawa, K., Vibe-Pederson, K. & Baralle, F. E. (1985) Primary structure of human fibronectin: differential splicing may generate at least 10 polypeptides from a single gene *EMBO J.* 4, 1755-1759.
- Kraulis, P. (1991) MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures. *J. Appl. Cryst.* 24, 946-950.
- Kunkel, T. A. (1985) Rapid and efficient site-specific mutagenesis without phenotypic selection. *Proc. Natl. Acad. Sci. USA* 82, 488-492.
- Leahy, D. J., Hendrickson, W. A., Aukhil, I. & Erickson, H. P. (1992) Structure of a fibronectin type III domain from tenascin phased by MAD analysis of the selenomethionyl protein *Science* 258, 987-991.
- Lee, W., Revington, M. J., Arrowsmith, C. & Kay, L. E. (1994) A pulsed field gradient isotope-filtered 3D ¹³C HMQC-NOESY experiment for extracting intermolecular NOE contacts in molecular complexes. *FEBS lett.* 350, 87-90.
- Lerner, R. A. & Barbas III, C. F., *Acta Chemica Scandinavica*, 50 672-678 (1996).
- Lesk, A. M. & Tramontano, A. (1992) Antibody structure and structural predictions useful in guiding antibody engineering. In *Antibody engineering. A practical guide*. (Borrebaeck, C. A. K., Ed.) Vol. W. H. Freeman & Co., New York.
- Li, B., Tom, J. Y., Oare, D., Yen, R., Fairbrother, W. J., Wells, J. A. & Cunningham, B. C. (1995) Minimization of a polypeptide hormone *Science* 270, 1657-1660.
- Litvinovich, S. V., Novokhatny, V. V., Brew, S. A. & Ingham, K. C. (1992) Reversible unfolding of an isolated heparin and DNA binding fragment, the first type III module from fibronectin. *Biochim. Biophys. Acta* 1119, 57-62.

- Logan, T. M., Olejniczak, E. T., Xu, R. X. & Fesik, S. W. (1992) Side chain and backbone assignments in isotopically labeled proteins from two heteronuclear triple resonance experiments. *FEBS lett.* 314, 413-418.
- Main, A. L., Harvey, T. S., Baron, M., Boyd, J. & Campbell, I. D. (1992) The three-dimensional structure of the tenth type III module of fibronectin: an insight into RGD-mediated interactions. *Cell* 71, 671-678.
- Masat, L., et al., (1994) *PNAS* 91:893-896.
- Martin, F., Toniatti, C., Ciliberto, G., Cortese, R. & Sollazzo, M. (1994) The affinity-selection of a minibody polypeptide inhibitor of human interleukin-6. *EMBO J.* 13, 5303-5309.
- Martin, M. T., *Drug Discov. Today*, 1:239-247 (1996).
- McCafferty, J., Griffiths, A. D., Winter, G. & Chiswell, D. J. (1990) Phage antibodies: filamentous phage displaying antibody variable domains. *Nature* 348, 552-554.
- McConnell, S. J., & Hoess, R. H., *J.Mol. Biol.* 250:460-470 (1995).
- Metzler, W. J., Leiting, B., Pryor, K., Mueller, L. & Farmer, B. T. I. (1996) The three-dimensional solution structure of the SH2 domain from p55blk kinase. *Biochemistry* 35, 6201-6211.
- Minor, D. L. J. & Kim, P. S. (1994) Measurement of the β -sheet-forming propensities of amino acids. *Nature* 367, 660-663.
- Muhandiram, D. R., Xu, G. Y. & Kay, L. E. (1993) An enhanced-sensitivity pure absorption gradient 4D ^{15}N , ^{13}C -edited NOESY experiment. *J. Biomol. NMR* 3, 463-470.
- Müller, C. W., Rey, F. A., Sodeoka, M., Verdine, G. L. & Harrison, S. C. (1995) Structure of the NH-kB p50 homodimer bound to DNA. *Nature* 373, 311-317.
- Nilges, M., Clore, G. M. & Gronenborn, A. M. (1988) Determination of three-dimensional structures of proteins from interproton distance data by hybrid

distance geometry-dynamical simulated annealing calculations. *FEBS lett.* 229, 317-324.

Nilges, M., Kuszewski, J. & Brünger, A. T. (1991) in *Computational aspects of the study of biological macromolecules by nuclear magnetic resonance spectroscopy*. (Hoch, J. C., Poulsen, F. M. and Redfield, C., Ed.) Vol. pp. 451-455, Plenum Press, New York.

O'Neil et al., (1994) in *Techniques in Protein Chemistry V* (Crabb, L., ed.) pp.517-524, Academic Press, San Diego.

O'Neil, K. T. & Hoess, R. H. (1995) Phage display: protein engineering by directed evolution. *Curr. Opinion Struct. Biol.* 5, 443-449.

Pace, C. N., Shirley, B. A. & Thomson, J. A. (1989) Measuring the conformational stability of a protein. In *Protein structure. A practical approach* (Creighton, T. E., Ed.) Vol. pp 311-330, IRL Press, Oxford.

Parmley, S. F. & Smith, G. P. (1988) Antibody-selectable filamentous fd phage vectors: affinity purification of target genes *Gene* 73, 305-318.

Pascal, S. M., Muhandiram, D. R., Yamazaki, T., Forman-Kay, J. D. & Kay, L. E. (1994a) Simultaneous acquisition of ¹⁵N- and ¹³C-edited NOE spectra of proteins dissolved in H₂O. *J. Magn. Reson. B* 103, 197-201.

Pascal, S. M., Singer, A. U., Gish, G., Yamazaki, T., Shoelson, S. E., Pawson, T., Kay, L. E. & Forman-Kay, J. D. (1994b) Nuclear magnetic resonance structure of an SH2 domain of phospholipase C-g1 complexed with a high affinity binding peptide. *Cell* 77, 461-472.

Pessi, A., Bianchi, E., Crameri, A., Venturini, S., Tramontano, A. & Sollazzo, M. (1993) A designed metal-binding protein with a novel fold. *Nature* 362, 3678-369.

Rees, A. R., Staunton, D., Webster, D. M., Searle, S. J., Henry, A. H. & Pedersen, J. T. (1994) Antibody design: beyond the natural limits. *Trends Biotechnol.* 12, 199-206.

Roberts et al., (1992) *Proc. Natl. Acad. Sci. USA* 89, 2429-2433.

- Rosenblum, J. S. & Barbas, C. F. I. (1995) in *Antobody Engineering* (Borrenbaeck, C. A. K., Ed.) Vol. pp89-116, Oxford University Press, Oxford.
- Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A laboratory manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor.
- Sandhu, G. S., Aleff, R. A. & Kline, B. C. (1992) Dual asymmetric PCR: one-step construction of synthetic genes. *BioTech.* 12, 14-16.
- Santoro, M. M. & Bolen, D. W. (1988) Unfolding free energy changes determined by the linear extrapolation method. 1. Unfolding of phenylmethanesulfonyl α -chymotrypsin using different denaturants *Biochemistry* 27, 8063-8068.
- Smith, G. P. & Scott, J. K. (1993) Libraries of peptides and proteins displayed on filamentous phage. *Methods Enzymol.* 217, 228-257.
- Smith, G. P. (1985) Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science* 228, 1315-1317.
- Smith, C. K., Munson, M. & Regan, L. (1995). Studying α -helix and β -sheet formation in small proteins. *Techniques Prot. Chem.*, 6, 323-332.
- Smith, C. K., Withka, J. M. & Regan, L. (1994) A thermodynamic scale for the β -sheet forming tendencies of the amino acids. *Biochemistry* 33, 5510-5517.
- Smyth, M. L. & von Itzstein, M. (1994) Design and synthesis of a biologically active antibody mimic based on an antibody-antigen crystal structure. *J. Am. Chem. Soc.* 116, 2725-2733.
- Studier, F. W., Rosenberg, A. H., Dunn, J. J. & Dubendorff, J. W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes *Methods Enzymol.* 185, 60-89.
- Suzuki, H. (1994) Recent advances in abzyme studies. *J. Biochem.* 115, 623-628.
- Tello, D., Goldbaum, F. A., Mariuzza, R. A., Ysern, X., Schwarz, F. P. & Poljak, R. J. (1993) Immunoglobulin superfamily interactions. *Biochem. Soc. Trans.* 21, 943-946.
- Thomas, N. R. (1994) Hapten design for the generation of catalytic antibodies. *Appl. Biochem. Biotech.* 47, 345-372.

- Verhoeyen, M., Milstein, C. & Winter, G. (1988) Reshaping human antibodies: Grafting an antilysozyme activity. *Science* 239, 1534-1536.
- Venturini et al., (1994) *Protein Peptide Letters* 1, 70-75.
- Vuister, G. W. & Bax, A. (1992) Resolution enhancement and spectral editing of uniformly ^{13}C -enriched proteins by homonuclear broadband ^{13}C decoupling. *J. Magn. Reson.* 98, 428-435.
- Vuister, G. W., Clore, G. M., Gronenborn, A. M., Powers, R., Garrett, D. S., Tschudin, R. & Bax, A. (1993) Increased resolution and improved spectral quality in four-dimensional $^{13}\text{C}/^{13}\text{C}$ -separated HMQC-NOESY-HMQC spectra using pulsed field gradients. *J. Magn. Reson. B* 101, 210-213.
- Ward, E. S., Güssow, D., Griffiths, A. D., Jones, P. T. & Winter, G. (1989) Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli* *Nature* 341, 554-546.
- Webster, D. M., Henry, A. H. & Rees, A. R. (1994) Antibody-antigen interactions *Curr. Opinion Struct. Biol.* 4, 123-129.
- Williams, A. F. & Barclay, A. N., *Ann. Rev. Immunol.* 6:381-405 (1988).
- Wilson, I. A. & Stanfield, R. L. (1993) Antibody-antigen interactions. *Curr. Opinion Struct. Biol.* 3, 113-118.
- Wilson, I. A. & Stanfield, R. L. (1994) Antibody-antigen interactions: new structures and new conformational changes *Curr. Opinion Struct Biol.* 4, 857-867.
- Winter, G., Griffiths, A. D., Hawkins, R. E. & Hoogenboom, H. R. (1994) Making antibodies by phage display technology *Annu. Rev. Immunol.* 12, 433-455.
- Wiseman, T., Williston, S., Brandts, J. F. & Lin, L.-N. (1989) Rapid measurement of binding constants and heats of binding using a new titration calorimeter. *Anal. Biolchem.* 179, 131-137.
- Wittenkind, M. & Mueller, L. (1993) HNCACB, a high-sensitivity 3D NMR experiment to correlate amide-proton and nitrogen resonances with the

- alpha- and beta-carbon resonances in proteins *J. Magn. Reson. B* 101, 201-205.
- Wu, T. T., Johnson, G. & Kabat, E. A. (1993) Length distribution of CDRH3 in antibodies *Proteins: Struct. Funct. Genet.* 16, 1-7.
- Wüthrich, K. (1986) NMR of proteins and nucleic acids, John Wiley & Sons, New York.
- Yamazaki, T., Forman-Kay, J. D. & Kay, L. E. (1993) Two-Dimensional NMR Experiments for Correlating ^{13}C -beta and ^1H -delta/epsilon Chemical Shifts of Aromatic Residues in ^{13}C -Labeled Proteins via Scalar Couplings. *J. Am. Chem. Soc.* 115, 11054.
- Zhang, O., Kay, L. E., Olivier, J. P. & Forman-Kay, J. D. (1994) Backbone ^1H and ^{15}N resonance assignments of the N-terminal SH3 domain of drk in folded and unfolded states using enhanced-sensitivity pulsed field gradient NMR techniques. *J. Biomol. NMR* 4, 845-858.

WHAT IS CLAIMED IS:

1. A fibronectin type III (Fn3) polypeptide monobody comprising a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences,
wherein one or more of the monobody loop region sequences vary by deletion, insertion or replacement of at least two amino acids from the corresponding loop region sequences in wild-type Fn3, and
wherein the β -strand domains of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequence of wild-type Fn3's β -strand domain sequences.
2. The monobody of claim 1, wherein at least one loop region is capable of binding to a specific binding partner (SBP) to form a polypeptide:SBP complex having a dissociation constant of less than 10^{-6} moles/liter.
3. The monobody of claim 1, wherein at least one loop region is capable of catalyzing a chemical reaction with a catalyzed rate constant (k_{cat}) and an uncatalyzed rate constant (k_{uncat}) such that the ratio of k_{cat}/k_{uncat} is greater than 10.
4. The monobody of claim 1, wherein one or more of the loop regions comprise amino acid residues:
 - i) from 15 to 16 inclusive in an AB loop;
 - ii) from 22 to 30 inclusive in a BC loop;
 - iii) from 39 to 45 inclusive in a CD loop;
 - iv) from 51 to 55 inclusive in a DE loop;
 - v) from 60 to 66 inclusive in an EF loop; and
 - vi) from 76 to 87 inclusive in an FG loop.

5. The monobody of claim 1, wherein the monobody loop region sequences vary from the wild-type Fn3 loop region sequences by the deletion or replacement of at least 2 amino acids.
6. The monobody of claim 1, wherein the monobody loop region sequences vary from the wild-type Fn3 loop region sequences by the insertion of from 3 to 25 amino acids.
7. An isolated nucleic acid molecule encoding the polypeptide monobody of claim 1.
8. An expression vector comprising an expression cassette operably linked to the nucleic acid molecule of claim 7.
9. The expression vector of claim 8, wherein the expression vector is an M13 phage-based plasmid.
10. A host cell comprising the vector of claim 8.
11. A method of preparing a fibronectin type III (Fn3) polypeptide monobody comprising the steps of:
 - a) providing a DNA sequence encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences wherein at least one loop region contains a unique restriction enzyme site;
 - b) cleaving the DNA sequence at the unique restriction site;
 - c) inserting into the restriction site a DNA segment known to encode a peptide capable of binding to a specific binding partner (SBP) or a

- transition state analog compound (TSAC) so as to yield a DNA molecule comprising the insertion and the DNA sequence of (a); and
- d) expressing the DNA molecule so as to yield polypeptide monobody.

12. A method of preparing a fibronectin type III (Fn3) polypeptide monobody comprising the steps of:
- (a) providing a replicatable DNA sequence encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein the nucleotide sequence of at least one loop region is known;
 - (b) preparing polymerase chain reaction (PCR) primers sufficiently complementary to the known loop sequence so as to be hybridizable under PCR conditions, wherein at least one of the primers contains a modified nucleic acid sequence to be inserted into the DNA;
 - (c) performing polymerase chain reaction using the DNA sequence of (a) and the primers of (b);
 - (d) annealing and extending the reaction products of (c) so as to yield a DNA product; and
 - (e) expressing the polypeptide monobody encoded by the DNA product of (d).
13. A method of preparing a fibronectin type III (Fn3) polypeptide monobody comprising the steps of:
- a) providing a replicatable DNA sequence encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein the nucleotide sequence of at least one loop region is known;
 - b) performing site-directed mutagenesis of at least one loop region so as to create a DNA sequence comprising an insertion mutation; and

- c) expressing the polypeptide monobody encoded by the DNA sequence comprising the insertion mutation.

14. A kit for performing the method of any one of claims 11-13, comprising a replicatable DNA encoding a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences.
15. A variegated nucleic acid library encoding Fn3 polypeptide monobodies comprising a plurality of nucleic acid species each comprising a plurality of loop regions, wherein the species encode a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences,
wherein one or more of the loop region sequences vary by deletion, insertion or replacement of at least two amino acids from corresponding loop region sequences in wild-type Fn3, and
wherein the β -strand domain sequences of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequences of β -strand domain sequences of the wild-type Fn3.
16. The variegated nucleic acid library of claim 15, wherein one or more of the loop regions encodes:
- i) an AB amino acid loop from residue 15 to 16 inclusive;
 - ii) a BC amino acid loop from residue 22 to 30 inclusive;
 - iii) a CD amino acid loop from residue 39 to 45 inclusive;
 - iv) a DE amino acid loop from residue 51 to 55 inclusive;
 - v) an EF amino acid loop from residue 60 to 66 inclusive; and
 - vi) an FG amino acid loop from residue 76 to 87 inclusive.

17. The variegated nucleic acid library of claim 15, wherein the loop region sequences vary from the wild-type Fn3 loop region sequences by the deletion or replacement of at least 2 amino acids.
18. The variegated nucleic acid library of claim 15, wherein the monobody loop region sequences vary from the wild-type Fn3 loop region sequences by the insertion of from 3 to 25 amino acids.
19. The variegated nucleic acid library of claim 15, wherein a variegated nucleic acid sequence comprising from 6 to 75 nucleic acid bases is inserted in any one of the loop regions of said species.
20. The variegated nucleic acid library of claim 15, wherein the variegated sequence is constructed so as to avoid one or more codons selected from the group consisting of those codons encoding cysteine or the stop codon.
21. The variegated nucleic acid library of claim 15, wherein the variegated nucleic acid sequence is located in the BC loop.
22. The variegated nucleic acid library of claim 15, wherein the variegated nucleic acid sequence is located in the DE loop.
23. The variegated nucleic acid library of claim 15, wherein the variegated nucleic acid sequence is located in the FG loop.
24. The variegated nucleic acid library of claim 15, wherein the variegated nucleic acid sequence is located in the AB loop.

25. The variegated nucleic acid library of claim 15, wherein the variegated nucleic acid sequence is located in the CD loop.
26. The variegated nucleic acid library of claim 15, wherein the variegated nucleic acid sequence is located in the EF loop.
27. A peptide display library derived from the variegated nucleic acid library of claim 15.
28. A peptide display library of claim 27, wherein the peptide is displayed on the surface of a bacteriophage or virus.
29. A peptide display library of claim 28, wherein the bacteriophage is M13 or fd.
30. A method of identifying the amino acid sequence of a polypeptide molecule capable of binding to a specific binding partner (SBP) so as to form a polypeptide:SSP complex wherein the dissociation constant of the said polypeptide:SBP complex is less than 10^{-6} moles/liter, comprising the steps of:
- providing a peptide display library according to claim 28;
 - contacting the peptide display library of (a) with an immobilized or separable SBP;
 - separating the peptide:SBP complexes from the free peptides,
 - causing the replication of the separated peptides of (c) so as to result in a new peptide display library distinguished from that in (a) by having a lowered diversity and by being enriched in displayed peptides capable of binding the SBP;

- e) optionally repeating steps (b), (c), and (d) with the new library of (d);
and
- f) determining the nucleic acid sequence of the region encoding the displayed peptide of a species from (d) and deducing the peptide sequence capable of binding to the SBP.

31. A method of preparing a variegated nucleic acid library encoding Fn3 polypeptide monobodies having a plurality of nucleic acid species each comprising a plurality of loop regions, wherein the species encode a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein one or more of the loop region sequences vary by deletion, insertion or replacement of at least two amino acids from corresponding loop region sequences in wild-type Fn3, and wherein the β -strand domain sequences of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequences of β -strand domain sequences of the wild-type Fn3, comprising the steps of
- a) preparing an Fn3 polypeptide monobody having a predetermined sequence;
 - b) contacting the polypeptide with a specific binding partner (SBP) so as to form a polypeptide:SSP complex wherein the dissociation constant of the said polypeptide:SBP complex is less than 10^{-6} moles/liter;
 - c) determining the binding structure of the polypeptide:SBP complex by nuclear magnetic resonance spectroscopy or X-ray crystallography;
and
 - d) preparing the variegated nucleic acid library, wherein the variegation is performed at positions in the nucleic acid sequence which, from the information provided in (c), result in one or more polypeptides with improved binding to the SBP.

32. A method of identifying the amino acid sequence of a polypeptide molecule capable of catalyzing a chemical reaction with a catalyzed rate constant, k_{cat} , and an uncatalyzed rate constant, k_{uncat} , such that the ratio of k_{cat}/k_{uncat} is greater than 10, comprising the steps of:
- a) providing a peptide display library according to claim 28;
 - b) contacting the peptide display library of (a) with an immobilized or separable transition state analog compound (TSAC) representing the approximate molecular transition state of the chemical reaction;
 - c) separating the peptide:TSAC complexes from the free peptides;
 - d) causing the replication of the separated peptides of (c) so as to result in a new peptide display library distinguished from that in (a) by having a lowered diversity and by being enriched in displayed peptides capable of binding the TSAC;
 - e) optionally repeating steps (b), (c), and (d) with the new library of (d); and
 - f) determining the nucleic acid sequence of the region encoding the displayed peptide of a species from (d) and hence deducing the peptide sequence.
33. A method of preparing a variegated nucleic acid library encoding Fn3 polypeptide monobodies having a plurality of nucleic acid species each comprising a plurality of loop regions, wherein the species encode a plurality of Fn3 β -strand domain sequences that are linked to a plurality of loop region sequences, wherein one or more of the loop region sequences vary by deletion, insertion or replacement of at least two amino acids from corresponding loop region sequences in wild-type Fn3, and wherein the β -strand domain sequences of the monobody have at least a 50% total amino acid sequence homology to the corresponding amino acid sequences of β -strand domain sequences of the wild-type Fn3, comprising the steps of

- a) preparing an Fn3 polypeptide monobody having a predetermined sequence, wherein the polypeptide is capable of catalyzing a chemical reaction with a catalyzed rate constant, k_{cat} , and an uncatalyzed rate constant, k_{uncat} , such that the ratio of k_{cat}/k_{uncat} is greater than 10;
- b) contacting the polypeptide with an immobilized or separable transition state analog compound (TSAC) representing the approximate molecular transition state of the chemical reaction;
- c) determining the binding structure of the polypeptide:TSAC complex by nuclear magnetic resonance spectroscopy or X-ray crystallography; and
- d) preparing the variegated nucleic acid library, wherein the variegation is performed at positions in the nucleic acid sequence which, from the information provided in (c), result in one or more polypeptides with improved binding to or stabilization of the TSAC.

34. An isolated polypeptide identified by the method of claim 30

35. An isolated polypeptide identified by the method of claim 32.

36. A kit for identifying the amino acid sequence of a polypeptide molecule capable of binding to a specific binding partner (SBP) so as to form a polypeptide:SSP complex wherein the dissociation constant of the said polypeptide:SBP complex is less than 10^{-6} moles/liter, comprising the peptide display library of claim 28.

37. A kit for identifying the amino acid sequence of a polypeptide molecule capable of catalyzing a chemical reaction with a catalyzed rate constant, k_{cat} ,

and an uncatalyzed rate constant, k_{uncat} , such that the ratio of $k_{\text{cat}}/k_{\text{uncat}}$ is greater than 10, comprising the peptide display library of claim 28.

38. A polypeptide derived by using the kit of claim 36.
39. A polypeptide derived by using the kit of claim 37.

ABSTRACT

A fibronectin type III (Fn3) polypeptide monobody, a nucleic acid molecule encoding said monobody, and a variegated nucleic acid library encoding said monobody, are provided by the invention. Also provided are methods of preparing a Fn3 polypeptide monobody, and kits to perform said methods. Further provided is a method of identifying the amino acid sequence of a polypeptide molecule capable of binding to a specific binding partner (SBP) so as to form a polypeptide:SSP complex, and a method of identifying the amino acid sequence of a polypeptide molecule capable of catalyzing a chemical reaction with a catalyzed rate constant, k_{cat} , and an uncatalyzed rate constant, k_{uncat} , such that the ratio of k_{cat}/k_{uncat} is greater than 10.

00120 472950

"Express Mail" mailing label number: EL709305975US
Date of Deposit: August 11, 2000
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231
Printed Name Shawn Hise
Signature [Signature]

"Express Mail" mailing label number: EM516503325US
Date of Deposit: June 12, 1998
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231
Printed Name Ann S. VIKSNINS
Signature [Signature]




FIGURE 1

[illegible]

FIGURE 3




FIGURE 4

NdeI
CATATGCAGGTTTCTGATGTTCCGCGTGACCTGGAAGTTGTTGCTGCGACCCCGACTAGC
MetGlnValSerAspValProArgAspLeuGluValValAlaAlaThrProThrSer
-2 -1 1 10

BclI PvuII PstI BsiWI
CTGCTGATCAGCTGGGATGCTCCTGCAGTTACCGTGCGTTATTACCGTATCACGTACGGT
LeuLeuIleSerTrpAspAlaProAlaValThrValArgTyrTyrArgIleThrTyrGly
20 30

EcoRI
GAAACCGGTGGTAACTCCCCGGTTCAGGAATTCACCTGTACCTGGTCCAAGTCTACTGCT
GluThrGlyGlyAsnSerProValGlnGluPheThrValProGlySerLysSerThrAla
40 50

SalI Bst1107I
ACCATCAGCGGCCTGAAACCGGGTGTCTGACTATACCATCACTGTATACGCTGTTACTGGC
ThrIleSerGlyLeuLysProGlyValAspTyrThrIleThrValTyrAlaValThrGly
60 70

SacI XhoI
CGTGGTGACAGCCCAGCGAGCTCCAAGCCAATCTCGATTAACTACCGTACCTAGTAACTC
ArgGlyAspSerProAlaSerSerLysProIleSerIleAsnTyrArgThr
80 90

BamHI
GAGGATCC

0963764-03100

FIGURE 5

00760-4492660




FIGURE 6




FIGURE 8




FIGURE 7

#1 10^{10}
others 5×10^9

ubiquitin-1





FIGURE 9



ubiquitin-2

FIGURE 10

007780-4792E950




FIGURE 11




FIGURE 12




FIGURE 13

09637614-081100




FIGURE 14




FIGURE 15




FIGURE 16

00760-1732E960




FIGURE 17a




FIGURE 17(b) & 17(c)

United States Patent Application

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled: Artificial Antibody Polypeptides.

The specification of which is attached hereto.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (see page 3 attached hereto).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on the basis of which priority is claimed:

No such applications have been filed.

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

Application Number
60/049,410

Filing Date
June 12, 1997

I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

No such applications have been filed.

I hereby appoint the following attorney(s) and/or patent agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith:

Anglin, J. Michael	Reg. No. 24,916	Forrest, Bradley A.	Reg. No. 30,837	Litman, Mark A.	Reg. No. 26,390
Arora, Sunee	Reg. No. P-42,267	Hale, Jeffrey D.	Reg. No. 40,012	Lundberg, Steven W.	Reg. No. 30,568
Bernkopf, Paul A.	Reg. No. P-41,615	Harris, Robert J.	Reg. No. 37,346	McCrackin, Ann M.	Reg. No. P-42,858
Bianchi, Timothy E.	Reg. No. 39,610	Hofmann, Rudolph P., Jr.	Reg. No. 38,187	Polglaze, Daniel J.	Reg. No. 39,801
Billion, Richard E.	Reg. No. 32,836	Holloway, Sheryl S.	Reg. No. 37,850	Provence, David L.	Reg. No. P-43,022
Brennan, Thomas F.	Reg. No. 35,075	Huchsch, Joseph C.	Reg. No. P-42,673	Schwegman, Michael L.	Reg. No. 25,816
Brooks, Edward J., III	Reg. No. 40,925	Klima-Silberg, Catherine I.	Reg. No. 40,052	Simboli, Paul B.	Reg. No. 38,616
Clark, Barbara J.	Reg. No. 38,107	Kluth, Daniel J.	Reg. No. 32,146	Slifer, Russell D.	Reg. No. 39,838
Drake, Eduardo E.	Reg. No. 40,594	Lacy, Rodney L.	Reg. No. 41,136	Terry, Kathleen R.	Reg. No. 31,884
Dryja, Michael A.	Reg. No. 39,662	Leffert, Thomas W.	Reg. No. 40,697	Viksnins, Ann S.	Reg. No. 37,748
Eubankson, Janet E.	Reg. No. 39,665	Lemaire, Charles A.	Reg. No. 36,198	Woessner, Warren D.	Reg. No. 30,440
Fogg, David N.	Reg. No. 35,138				

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/organization/who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Schwegman, Lundberg, Woessner & Kluth, P.A. to the contrary.

Please direct all correspondence in this case to Schwegman, Lundberg, Woessner & Kluth, P.A. at the address indicated below:

P.O. Box 2938, Minneapolis, MN 55402
Telephone No. (612)373-6900

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of sole inventor : Shohei Koide
Citizenship: Japan Residence: Rochester, NY
Post Office Address: 236 Oakdale Drive
Rochester, NY 14618

Signature: Shohei Koide Date: 6/11/98
Shohei Koide

Full Name of inventor:
Citizenship: Residence:
Post Office Address:

Signature: _____ Date: _____

Full Name of inventor:
Citizenship: Residence:
Post Office Address:

Signature: _____ Date: _____

Full Name of inventor:
Citizenship: Residence:
Post Office Address:

Signature: _____ Date: _____

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

- (1) prior art cited in search reports of a foreign patent office in a counterpart application, and
- (2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim; or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office, or
 - (ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

- (1) Each inventor named in the application;
- (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

S/N Unknown

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Shohei Koide

Examiner: Unknown

Serial No.: Unknown

Group Art Unit: Unknown

Filed: Herewith

Docket: 109.034US3

Title: ARTIFICIAL ANTIBODY POLYPEPTIDES

COMMUNICATION UNDER 37 C.F.R. § 1.821

BOX PATENT APPLICATION

Commissioner for Patents

Washington, D.C. 20231

Sir:

This Communication is submitted to conform the above-referenced application to the requirements of 37 C.F.R. §§ 1.821 through 1.825.

The paper copy of the Sequence Listing in this application is identical to the computer readable copy of the Sequence Listing filed in application Serial No. 09/096,749, filed June 12, 1998. In accordance with 37 C.F.R. § 1.821(e), please use the last-filed computer readable form in that application as the computer readable form for the instant application. A paper copy of the Sequence Listing is included in a separately-filed preliminary amendment for incorporation into the specification.

Respectfully submitted,

SHOHEI KOIDE

By his Representatives,

SCHWEGMAN, LUNDBERG, WOESSNER & KLUTH, P.A.

P.O. Box 2938

Minneapolis, MN 55402

(612) 373-6961

Date

11 August 2000

By



Ann S. Viksnins

Reg. No. 37,748

Express Mail No.: EL709305975US

Mailing Date: August 11, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Box PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.

Shawn L. Hise
Printed Name


Signature

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: Koide, Shohei
- (ii) TITLE OF THE INVENTION: ARTIFICIAL ANTIBODY POLYPEPTIDES
- (iii) NUMBER OF SEQUENCES: 118
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Schwegman, Lundberg, Woessner & Kluth P.A.
 - (B) STREET: 121 South Eighth Street, Ste. 1600
 - (C) CITY: Minneapolis
 - (D) STATE: MN
 - (E) COUNTRY: USA
 - (F) ZIP: 55402
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 2.0b
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER: Unknown
 - (B) FILING DATE: Herewith
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: 09/096,749
 - (B) FILING DATE: June 12, 1998
- (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Ann S. Viksnins
 - (B) REGISTRATION NUMBER: 37,748
 - (C) REFERENCE/DOCKET NUMBER: 109.034US3
- (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: (612) 373-6900
 - (B) TELEFAX: (612) 339-3061

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 14 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

00730-4792950

Ala Arg Glu Arg Asp Tyr Arg Leu Asp Tyr Trp Gly Gln Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ala Arg Gly Ala Val Val Ser Tyr Tyr Ala Met Asp Tyr Trp Gly Gln
1 5 10 15
Gly

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Tyr Ala Val Thr Gly Arg Gly Asp Ser Pro Ala Ser Ser Lys Pro Ile
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

1 5 10

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Tyr Ala Val Arg Asp Tyr Arg Ser Lys Pro Ile
1 5 10

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Tyr Ala Val Thr Arg Asp Tyr Arg Leu Ser Ser Lys Pro Ile
1 5 10

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Tyr Ala Val Thr Glu Arg Asp Tyr Arg Leu Ser Ser Lys Pro Ile
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Tyr Ala Val Ala Val Val Ser Tyr Tyr Ala Met Asp Tyr Pro Ile
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Tyr Ala Val Thr Ala Val Val Ser Tyr Tyr Ala Ser Ser Lys Pro Ile
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 59 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CGGGATCCCA TATGCAGGTT TCTGATGTTT CGCGTGACCT GGAAGTTGTT GCTGCGACC

59

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TAACTGCAGG AGCATCCCAG CTGATCAGCA GGCTAGTCGG GGTCGCAGCA ACAAC

55

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CTCCTGCAGT TACCGTGCCT TATTACCGTA TCACGTACGG TGAAACCGGT G

51

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GTGAATTCCT GAACCGGGGA GTTACCACCG GTTTCACCG

39

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AGGAATTCAC TGTACCTGGT TCCAAGTCTA CTGCTACCAT CAGCGG

46

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GTATAGTCGA CACCCGGTTT CAGGCCGCTG ATGGTAGC

38

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGGGTGTGCA CTATACCATC ACTGTATACG CT

32

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 55 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CGGGATCCGA GCTCGCTGGG CTGTCACCCAC GGCCAGTAAC AGCGTATACA GTGAT

55

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CAGCGAGCTC CAAGCCAATC TCGATTAACT ACCGT

35

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CGGGATCCTC GAGTTACTAG GTACGCTAGT TAATCGA

37

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CGGGATCCAC GCGTGCCACC GGTACGGTAG TTAATCGA

38

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 44 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CGGGATCCAC GCGTCCATTC GTTTGTGAAT ATCAAGGCCA ATCG

44

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CCGGAAGCTT TAAGACTCCT TATTACGCAG TATGTTAGC

39

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

CTGTTACTGG CCGTGAGATC TAACCAGCGA GCTCCA

36

00130-492550

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GATCAGCTGG GATGCTCCTN NKNNKNNKNN KNNKTATTAC CGTATCACGT A

51

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

TGTATACGCT GTTACTGGCN NKNNKNNKNN KNNKNNKNNK TCCAAGCCAA TCTCGAT

57

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CTGTATACGC TGTTACTGGC NNKNNKNNKN NKCCAGCGAG CTCCAAG

47

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 base pairs

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Cys Trp Arg Arg Ala
1 5

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Arg Trp Val Gly Leu Ala Trp
1 5

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE: internal
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Cys Lys His Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Phe Ala Asp Leu Trp Trp Arg
1 5

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Cys Arg Arg Gly Arg
1 5

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Arg Gly Phe Met Trp Leu Ser
1 5

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

[illegible]

Cys Asn Trp Arg Arg
1 5

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Arg Ala Tyr Arg Tyr Arg Trp
1 5

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Ser Arg Leu Arg Arg
1 5

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

14

(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Pro Pro Trp Arg Val
1 5

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Ala Arg Trp Thr Leu
1 5

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Arg Arg Trp Trp Trp
1 5

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

Gly Gln Arg Thr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Arg Arg Trp Trp Ala
1 5

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Ala Val Thr Val Arg
1 5

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

Arg Gly Asp Ser Pro Ala Ser
1 5

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

Cys Asn Trp Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

Arg Ala Tyr Arg Tyr Arg Trp
1 5

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

[illegible]

Cys Met Trp Arg Ala
1 5

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Arg Trp Gly Met Leu Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Ala Arg Met Arg Glu
1 5

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

Arg Trp Leu Arg Gly Arg Tyr
1 5

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Cys Ala Arg Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

Arg Arg Ala Gly Trp Gly Trp
1 5

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Cys Asn Trp Arg Arg

1

5

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Arg Ala Tyr Arg Tyr Arg Trp

1

5

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Arg Trp Arg Glu Arg

1

5

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Arg His Pro Trp Thr Glu Arg

1

5

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Cys Asn Trp Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Arg Ala Tyr Arg Tyr Arg Trp
1 5

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Glu Arg Arg Val Pro
1 5

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

Arg Leu Leu Leu Trp Gln Arg
 1 5

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

Gly Arg Gly Ala Gly
 1 5

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Phe Gly Ser Phe Glu Arg Arg
 1 5

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:

[illegible]

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

(2) INFORMATION FOR SEQ ID NO:68:

```
(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

(2) INFORMATION FOR SEQ ID NO:69:

```
(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

(2) INFORMATION FOR SEQ ID NO:70:

23

(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

Arg Ala Tyr Arg Tyr Arg Trp
1 5

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

Ala Val Thr Val Arg
1 5

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

Gly Arg Gly Asp Ser
1 5

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

Arg Arg Trp Trp Ala
1 5

(2) INFORMATION FOR SEQ ID NO: 77:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

Gly Gln Arg Thr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

Arg Arg Trp Trp Ala
1 5

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

Leu Arg Tyr Arg Ser
1 5

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

Gly Trp Arg Trp Arg
1 5

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

Gly Gln Arg Thr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

Arg Arg Trp Trp Ala
1 5

(2) INFORMATION FOR SEQ ID NO:83:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

Gly Gln Arg Thr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

Arg Arg Trp Trp Ala
1 5

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

Leu Arg Tyr Arg Ser
1 5

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

Gly Trp Arg Trp Arg
1 5

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

Leu Arg Tyr Arg Ser
1 5

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

Gly Trp Arg Trp Arg
1 5

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

Gly Gln Arg Thr Phe
1 5

(2) INFORMATION FOR SEQ ID NO:90:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

Arg Arg Trp Trp Ala
1 5

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

Ala Val Trp Val Arg
1 5

(2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

GGCCGTGGTG ACAGCCCAGC GAGC

24

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

Gly Arg Gly Asp Ser Pro Ala Ser
1 5

(2) INFORMATION FOR SEQ ID NO:97:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

(2) INFORMATION FOR SEQ ID NO:98:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

Ser Arg Leu Arg Arg

1

5

(2) INFORMATION FOR SEQ ID NO:99:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

CCGCCGTGGA GGGTG

15

(2) INFORMATION FOR SEQ ID NO:100:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

Pro Pro Trp Arg Val

1

5

(2) INFORMATION FOR SEQ ID NO:101:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

GGTCAGCGAA CTTTT

15

(2) INFORMATION FOR SEQ ID NO:102:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

Gly Gln Arg Trp Phe

1

5

(2) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

AGGCGGTGGT GGGCT

15

(2) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

AGGCGGTGGT GGTGG

15

(2) INFORMATION FOR SEQ ID NO:108:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

Arg Arg Trp Trp Trp
1 5

(2) INFORMATION FOR SEQ ID NO:109:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

Gly Lys Lys Gly Lys
1 5

(2) INFORMATION FOR SEQ ID NO:110:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE: internal
 (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala	Ala	Thr	Pro	Thr	Ser	Leu	Leu
1				5				10						15				
Ile	Ser	Trp	Asp	Ala	Pro	Ala	Val	Thr	Val	Arg	Tyr	Tyr	Arg	Ile	Thr	Tyr	Gly	Glu
20				25				30						35				
Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu	Phe	Thr	Val	Pro	Gly	Ser	Lys	Ser	Thr	Ala
40				45				50						55				
Thr	Ile	Ser	Gly	Leu	Lys	Pro	Gly	Val	Asp	Tyr	Thr	Ile	Thr	Val	Tyr	Ala	Val	Thr
	60			65				70						75				
Gly	Arg	Gly	Asp	Ser	Pro	Ala	Ser	Ser	Lys	Pro	Ile	Ser	Ile	Asn	Tyr	Arg	Thr	
	80			85				90										

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 308 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE:
 (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

CATATGCAGG	TTTCTGATGT	TCCGCGTGAC	CTGGAAGTTG	TTGCTGCGAC	CCCGACTAGC	60
CTGCTGATCA	GCTGGGATGC	TCCTGCAGTT	ACCGTGCGTT	ATTACCGTAT	CACGTACGGT	120
GAAACCGGTG	GTAAC TCCCC	GGTTCAGGAA	TTC ACTGTAC	CTGGTTCCAA	GTCTACTGCT	180
ACCATCAGCG	GCCTGAAACC	GGGTGTCGAC	TATACCATCA	CTGTATACGC	TGTTACTGGC	240
CGTGGTGACA	GCCCAGCGAG	CTCCAAGCCA	ATCTCGATTA	ACTACCGTAC	CTAGTAACTC	300
GAGGATCC						308

(2) INFORMATION FOR SEQ ID NO:112:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 96 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO
(v) FRAGMENT TYPE: internal
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

Met	Gln	Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala	Ala	Thr	Pro	Thr	Ser
1				5					10					15				
Leu	Leu	Ile	Ser	Trp	Asp	Ala	Pro	Ala	Val	Thr	Val	Arg	Tyr	Tyr	Arg	Ile	Thr	Tyr
20				25					30					35				
Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu	Phe	Thr	Val	Pro	Gly	Ser	Lys	Ser
40				45					50					55				
Thr	Ala	Thr	Ile	Ser	Gly	Leu	Lys	Pro	Gly	Val	Asp	Tyr	Thr	Ile	Thr	Val	Tyr	Ala
60				65					70					75				
Val	Thr	Gly	Arg	Gly	Asp	Ser	Pro	Ala	Ser	Ser	Lys	Pro	Ile	Ser	Ile	Asn	Tyr	Arg
80				85					90					95				

Thr

(2) INFORMATION FOR SEQ ID NO:113:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

Arg Gly Asp
1

(2) INFORMATION FOR SEQ ID NO:114:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

Met	Gly	Ser	Ser	His	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro
1				5				10				15		
Arg	Gly	Ser	His											
20														

(2) INFORMATION FOR SEQ ID NO:115:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

Gly Gln Arg Thr Phe Arg Arg Trp Trp Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:116:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

Leu Arg Tyr Arg Ser Gly Trp Arg Trp Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO:117:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

Cys Asn Trp Arg Arg Arg Ala Tyr Arg Tyr Trp Arg
1 5 10

(2) INFORMATION FOR SEQ ID NO:118:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

Ala	Arg	Met	Arg	Glu	Arg	Trp	Leu	Arg	Gly	Arg	Tyr
1				5					10		